

Formation Capital Corporation, U.S. 812 Shoup Street Salmon, ID 83467

Tel: 208.756.4578 Fax: 208.756.2573



June 14, 2013

US EPA Region 10 Attn: PCS Data Entry Team 1200 Sixth Avenue Suite 900, OCE-133 Seattle, WA 98101

Subject:

2012 Fish Tissue & Macro Invertebrate Monitoring Results

Formation Capital Corporation, U.S. (Formation), submits the attached 2012 Fish Tissue & Macro Invertebrate Monitoring Results for the Idaho Cobalt Project (ICP) pursuant to the NPDES Permit. A hard copy of this submittal is provided for your and those courtesy copied on this transmittal.

If you have any questions about this submittal please contact me by telephone at 208-756-4578 ext. 26 or by email at wgscales@formcap.com.

Respectfully,

Wm. G. (Bill) Scales

President

Attachment

CC:

IDEQ Attn: Water Division NMFS Attn: Director

U.S. Fish and Wildlife

I (IS)

IDAHO COBALT PROJECT

FISH TISSUE & MACROINVERTEBRATE MONITORING RESULTS

In accordance with
United States Environmental Protection Agency
National Pollutant Discharge Elimination System
Permit No. ID-002832-1

Prepared for:

Formation Capital Corporation, U.S. 812 Shoup Street Salmon, Idaho 83467

Prepared by:

Hildebrand

Hildebrand & Associates, LLC 4623 E. Pegasus Ct Boise, Idaho 83716

> December 11, 2012 (Revised January 3, 2013)

	CONTENTS	
<u>Secti</u>	<u>on</u>	Page
1.0	INTRODUCTION	2
2.0	SAMPLING APPROACH	
	2.1 SITE SELECTION AND LOCATIONS	2_1
3.0	SAMPLING METHODS	
	3.1 IN SITU WATER QUALITY	2-2
	3.2 MACROINVERTEBRATE SAMPLING	2-2
	3.3 MACROINVERTEBRATE HANDLING AND ANALYSIS	2-3
	3.4 BIOTIC INDICES	
	3.5 HABITAT ASSESSMENT	
	3.6 SUBSTRATE CHARACTERIZATION	2-5
	3.7 FISH TISSUE COLLECTION	2-5
	3.8 FISH TISSUE HANDLING AND ANALYSIS	2-6
	3.9 CHAIN OF CUSTODY	2-11
	3.9.1 QUALITY CONTROL AND QUALITY ASSURANCE	
4.0	RESULTS AND DISCUSSION	2-12
	4.1 IN SITU WATER QUALITY	2-12
	4.2 MACROINVERTEBRATE SAMPLE.RESULTS	
	4.3 HABITAT ASSESSMENT	
	4.4 STTREAM DISCHARGE	2-16
	4.5 SUBSTRATE CHARACTERIZATION	2-17
	4.6 FISH TISSUE	
	4.7 CONCLUSION	
5.0	REFERENCES	2-20
	APPENDICES	
A B C D E F	YSI CALIBRATION SCIENTIFIC COLLECTING PERMIT AND REPORTING DOCUMEN' FISH COLLECTION FORMS CHAIN OF CUSTODY FORMS MACROINVERTEBRATE RESULTS WOLMAN PEBBLE COUNT FORM	TS
G	FISH TISSUE METALS	

FIGURES

FIGURE 1	MACROINVERTEBRATE SAMPLING LOCATIONS
FIGURE 2	FISH TISSUE SAMPLING LOCATIONS - BIG DEER CREEK
FIGURE 3	FISH TISSUE SAMPLING LOATIONS - UPPER BLACKBIRD CREEK

TABLES

TABLE 1	IN SITU WATER QUALITY MEASUREMENTS	
TABLE 2	PHYSICAL HABITAT DATA	
TABLE 3	STREAM DISCHARGE MEASUREMENTS	
TABLE 4	BIG DEER CREEK 01 FISH TISSUE METALS ANALYSIS (DRY WEIGH	HT)
TABLE 5	BIG DEER CREEK 01 FISH TISSUE METALS ANALYSIS (WET WEIG	HT)
TABLE 6	BIG DEER CREEK 02 FISH TISSUE METALS ANALYSIS (DRY WEIG	HT)
TABLE 7	BIG DEER CREEK 02 FISH TISSUE METALS ANALYSIS (WET WEIG	HT)
TABLE 8	UPPER BLACKBIRD CREEK FISH TISSUE METALS ANALYSIS (DRY	WEIGHT)
TABLE 9	UPPER BLACKBIRD CREEK FISH TISSUE METALS ANALYSIS (WET	WEIGHT)
TABLE 10	METHOD BLANKS DRY WEIGHT BASIS	3
TABLE II	METHOD BLANKS WET WEIGHT BASIS	
TABLE 12	MERCURY METHOD BLANKS DRY WEIGHT BASIS	19
TABLE 13	MERCURY METHOD BLANKS WET WEIGHT BASIS	

1.0 INTRODUCTION

Formation Capital Corporation, U.S. (Formation) retained Hildebrand & Associates, LLC (Hildebrand) to collect aquatic benthic macroinvertebrates (Macroinvertebrates) and fish tissue samples to evaluate the macroinvertebrate community structure and bioaccumulation of trace metals in fish tissue in Big Deer Creek and in Upper Blackbird Creek located within Salmon-Cobalt Ranger District of the Salmon National Forest. This biological monitoring project and summary report has been prepared for Formation's Idaho Cobalt Project in accordance with Formation's final National Pollutant Discharge Elimination System (NPDES) permit number ID-002832-1.

2.0 SAMPLING APPROACH

Based on locations identified in the Idaho Cobalt NPDES Quality Assurance Project Plan (TELESTO, 2010) and Biological Monitoring and Assessment Program (ENVIRON, 2009), one macroinvertebrate monitoring station and three fish tissue monitoring stations were established within the Salmon-Cobalt Ranger District of the Salmon National Forest. Samples were collected from September 4th, 2012 through September 5th, 2012.

2.1 SITE SELECTION AND LOCATIONS

One macroinvertebrate monitoring station (BDC01) and one fish tissue monitoring station (BDC-01-01) was established in Big Deer Creek, downstream of Formation's proposed water treatment discharge outfall 001. Site BDC01 was selected to assess potential changes in the macroinvertebrate community structure. Site BDC-01-01 was selected to assess potential fish tissue bioaccumulation. Fish tissue concentrations are monitored for bioaccumulation of metals so that tissue concentrations can be compared prior to discharge to post discharge. Macroinvertebrate community structure is monitored so that macroinvertebrate composition can be compared prior to discharge to post discharge.

One fish tissue monitoring station was established in Big Deer Creek (BDC-02) upstream of Formation's proposed water treatment discharge outfall 001. An additional fish tissue monitoring reference station was established in Upper Blackbird Creek (UBC-01). The macroinvertebrate sampling locations are presented in Figure 1. Fish tissue monitoring locations are presented in Figure 2 and Figure 3.

3.0 SAMPLING METHODS

Beneficial Use Reconnaissance Program Protocols (BURP) developed by the Idaho Department of Environmental Quality (DEQ, 2007) was used to evaluate the biological assemblages and physical habitat structure. Included in the BURP are measures of water quality, collection of benthic macroinveretbrates and evaluations of physical habitat. Embeddedness and substrate size, one of the most important determinations of habitat for fish and macroinvertebrates were evaluated by conducting a modified Wolman Pebble Count (Wolman et al., 1954) to quantify substrate size distribution in riffle habitats. Sampling methods are described below.

3.1 IN SITU WATER QUALITY

In situ water quality measurements were collected at each monitoring station including the freshwater fish pond and included instantaneous measurements of pH (standard units), conductivity (micromhos per centimeter) [µmhos/cm]), water temperature (degrees Celsius [°C]), dissolved oxygen (DO) (milligrams per liter [mg/L]) and percent DO saturation. These measurements serve to identify water quality conditions which may affect aquatic life. In situ water quality measurements were collected prior to collection of fish and macroinvertebrates and habitat evaluations. An YSI 556 multi-parameter field instrument, calibrated prior to use, was positioned at approximately 0.5 feet in the water column in an undisturbed area of the monitoring station. Once the YSI stabilized, in situ water quality measurements were recorded. YSI 556 calibration results are included in Appendix A. In addition, stream discharge was measured utilizing a Marsh McBirney Flo-Mate 2000 Digital Flow Meter.

3.2 MACROINVERTEBRATE SAMPLING

Macroinvertebrates were collected on September 4th, 2012 under Idaho Department of Fish and Game (IDFG) Scientific Collecting Permit No. F-12-12-12 (Appendix B). To minimize impacts to the benthic community in Big Deer Creek, macroinvertebrate community sampling was only performed. Due to their limited mobility and relatively long life span, macroinvertebrates integrate and reflect water quality effects over time and are excellent indicators of stress in aquatic systems. Additionally, macroinvertebrates with certain environmental tolerances may provide some insight into the presence of water toxicity.

Macroinvertebrate sampling followed bioassessment methodologies of the IDEQ BURP. The IDEQ BURP methodology includes qualitative habitat evaluations, macroinvertebrate collections and processing techniques, and taxa identification procedures. For the Idaho Cobalt Project, quantitative macroinvertebrate samples were collected using a Hess Sampler fitted with a 500 micron (μm) mesh collection net. The macroinvertebrate sampling reach was approximately 30-times the bankfull width which equated to 450 feet (137 meters). Three evenly spaced riffle transects (T1, T2, and T3) (Figure 1) were established within BDC01and macroinvertebrate samples were collected from each separate transect for a total of three macroinvertebrate samples. To provide consistency in the sampling design, the Hess Sampler was used to collect macroinvertebrates at all 3 transects.

3.3 MACROINVERTEBRATE HANDLING AND ANALYSIS

Macroinvertebrate samples were stored and preserved in individual 1-liter Nalgene® High-Density Polyethylene (HDPE) sample containers with 99 percent isopropyl. Sample containers were labeled both inside and outside with labels containing the following information: station number, stream name, date and time of collection, and sample type. Macroinvertebrate samples were checked for adequate preservation, placed inside sampling coolers, and were secured in locked field vehicles for the duration of the project.

Upon return from the field, macroinvertebrate samples were shipped by ground transport to EcoAnalysts, Inc., in Moscow, Idaho. Marcoinvertebrate samples were processed following the IDEQ BURP methodology of removing the first 500 animals and identifying the invertebrates to species or genus/species subgroups whenever possible. Each sample was mixed thoroughly on a sieve and split into eight homogeneous fractions. Each of the eight fractions was placed in a numbered tray. A random number generator was used to select the order in which the trays were sorted until the required total of 500 organisms per sample was reached or, if there were fewer than 500 organisms present in a sample, until all organisms were removed. The fraction of sample sorted to obtain the 500 invertebrates was recorded so that invertebrate abundance can be converted to number of organisms/meter squared (m²) for comparison to future studies.

3.4 BIOTIC INDICES

Macroinvertebrate data was evaluated using the Stream Macroinvertebrate Index (SMI) developed by the IDEQ as part of their ecological assessment approach to determine appropriate life use support in Idaho's streams. The component metrics of the SMI are as follows:

- Total Taxa Number of distinct taxa found in the macroinvertebrate assemblage at each station. Generally, the number of taxa decreases in response to increasing perturbation.
- Ephemeroptera Taxa Number of mayfly taxa, which generally decreases in response to increasing perturbation.
- Plecoptera Taxa Number of stonefly taxa, which generally decreases in response to increasing perturbation.
- Trichoptera Taxa Number of caddisfly taxa, which generally decreases in response to increasing perturbation.
- % Plecoptera Percent of sample that is stonefly nymphs. Predicted to decrease in response to increasing perturbation.
- Hilsenhoff Biotic Index (HBI) Abundance-weighted average tolerance to pollution. Originally a
 measure of tolerance to organic pollutants, but commonly used for evaluating responses to
 organic and toxic pollutants. Predicted to increase in response to increasing perturbation.
- % 5 Dominant Taxa Percent of the sample that are the five most dominant taxa. Predicted to increase in response to increasing perturbation.
- Scraper Taxa Number of invertebrate taxa that feed by scraping. This metric provides an
 indication of the riffle community food base (e.g. periphyton). Scrapers increase with increased
 abundance of periphyton and decrease as fine particle material increases. Their abundance
 generally decreases in response to increasing perturbation.
- Clinger Taxa Number of invertebrate taxa that are clingers. Clingers have fixed retreats or
 adaptations for attaching to surfaces in flowing water. They are adapted to life in running waters
 and are sensitive to hydrologic perturbation, habitat disturbance, and other pollutants. Their
 abundance generally decreases in response to increasing perturbation.

In addition to the metrics used in the calculation of the SMI, total invertebrate abundance (density), mayfly abundance, % mayflies, biomass and an index of metal tolerance was evaluated.

3.5 HABITAT ASESSMENT

Physical habitat quality is a major determinant of biological diversity of stream macroinvertebrate communities. In conjunction with macroinvertebrate sampling, a visual qualitative aquatic and riparian habitat assessment was completed in BDC - 01 and BDC - 02. The visual assessment of aquatic and riparian habitat is consistent with IDEQ BURP and parameters assessed as part of the habitat evaluation include stream discharge, canopy cover, in-stream cover, embeddedness in riffles and pools, channel shape, disruptive pressure, zone of influence, pool substrate characteristics and pool variability. Additional stream characteristics were recorded and included large organic debris, pool count, stream sinuosity, and Rosgen stream type. While these additional stream characteristics do not receive habitat assessment scores they still provide valuable stream habitat data that can be evaluated through time.

3.6 SUBSTRATE SIZE CHARACTERIZATION

Substrate characteristics are important determinants of habitat for fish and macroinvertebrates in streams (Kaufmann and Robison 1998, Kaufmann et al. 1999), and are often sensitive indicators of anthropogenic impacts on streams (Minshall et al 1985). Substrate size characterization was used to evaluate pre and post discharge effects on BDC - 01. Cobble-sized substrate provides the greatest amount of usable habitat to benthic macroinvertebrates, while smaller sized substrate offers reduced habitat for colonization (Green et al. 2000). Substrate size characterization was evaluated using the modified Wolman Pebble Count (Wolman et al., 1954). Results of the pebble count are discussed in Section 4.5.

3.7 FISH TISSUE COLLECTION

A total of 30 fish tissue samples were collected by rod and reel during September 4th and 5th, 2012 under IDFG Scientific Collecting Permit No. F-12-12-12 (Appendix B). Fish tissue samples were collected following the protocols as found in IDEQ's Implementation Guidance for the Idaho Mercury Water Quality Criteria (IDEQ 2005).

Ten individual fish tissue samples were collected from each monitoring station (Figure 2 and Figure 3). The species along with species length, sample location, date and time of collection and GPS coordinates were recorded on standardized field data sheets are provided in Appendix C. Each collected fish was also digitally photographed with an individual sample identifier based on the date and sample location. Digital photographs are provided on compact disk. Once collected and logged fish tissue samples were packaged and stored on ice in a cooler for the duration of the sampling event. Samples were frozen upon return


from the field and then were shipped on dry ice to ALS Environmental (formerly Columbia Analytical Services) in Kelso, Washington for chemical analyses.

3.8 FISH TISSUE HANDLING AND ANALYSIS


Fish tissue samples were handled according to the protocols outlined in the IDEQ's Implementation Guidance for the Idaho Mercury Water Quality Criteria (IDEQ 2005) which were adopted from the Idaho Fish Consumption Advisory Program (IFCAP) and United States Geological Survey (USGS) sampling techniques. This included clean sampling handling techniques for low level metals analysis to preclude false positives arising from sample collection, sample handling, or analysis.

Whole individual fish samples were wrapped in Glad ® Plastic Wrap and packaged individually in resealable plastic bags (e.g. Ziploc™). To ensure sample bags did not leak, samples were individually double bagged and samples from an individual site were packaged in a single, larger Ziploc™ bag. To ensure tissue samples were prepared in a laboratory clean-room environment using non-contaminating techniques, whole fish samples were shipped to ALS Environmental in Kelso, Washington for chemical analysis. Per the IDEQ mercury water quality criteria guidance (IDEQ, 2005), all fish tissue samples were analyzed as fillets without skin. Since the proposed sample number is relatively small, individual fillet samples were homogenized and were analyzed for total aluminum, arsenic, cadmium, cobalt, lead, manganese, mercury, nickel selenium, thallium and zinc. Fish tissue samples were first analyzed for total mercury and would be only analyzed for methylmercury if total mercury concentrations were within 20 percent of the IDEQ fish tissue mercury criterion of 0.3 milligrams per kilogram on a wet weight basis (mg/kg ww).


FORMATION CAPITAL CORPORATION, U.S. IDAHO COBALT PROJECT NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT



FORMATION CAPITAL CORPORATION, U.S. IDAHO COBALT PROJECT NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT



FORMATION CAPITAL CORPORATION, U.S. IDAHO COBALT PROJECT NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT



THIS PAGE LEFT INTENTIONALLY BLANK

FORMATION CAPITAL CORPORATION, U.S. IDAHO COBALT PROJECT NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

3.9 CHAIN OF CUSTODY

Samples collected in the field represent physical evidence and their possession must be traceable from the time of collection until the data are ultimately used. Proper chain of custody (COC) procedures was used to maintain and document sample possession. Completed COC forms accompanied each macroinvertebrate and fish tissue sample to their respective laboratory. Executed COC forms are provided in Appendix D.

3.9.1 QUALITY ASSURANCE AND QUALITY CONTROL

An inherent difficulty exists in generating blanks for solid samples such as fish; there is no blank tissue that can be used. Fish fillets are inherently difficult to contaminate because they are a solid medium. Only the surface is exposed to contamination before being pureed. Also, the much higher concentrations of mercury and other contaminants in fillets compared to water are not easily changed by small amounts of contamination. Since there will be no dissecting or filleting of fish in the field, deionized water washes or rinses of equipment did not take place. Therefore, field blanks were not analyzed in this study.

Homogenization blanks were used to assess potential contamination from analytical equipment. Homogenization blanks are equivalent to equipment blanks and were collected by running deionized water through the homogenization equipment. ALS Environmental analyzed one homogenization blank per sampling round or 1 for every 10 samples, whichever is greater. Method blanks were also used to assess potential contamination from analytical equipment. ALS Environmental analyzed one method blank for every sample batch or every 20 samples, whichever is more frequent. Method blanks consist of an analyte-free matrix that is subjected to the entire analytical process.

For analyzing macroinvertebrate samples, performance objectives associated primarily with measurement error were followed. Measurement error and performance objectives followed EPA Guidance for Quality Assurance Plans EPA240/R-02/009.

4.0 RESULTS AND DISCUSSION

4.1 IN SITU WATER QUALITY

In situ water quality measurements (pH, conductivity, water temperature, dissolved oxygen (DO) and DO percent saturation) were collected at each of the fish tissue and macroinvertebrate sampling stations including the freshwater fish pond (Table 1). The pH values are not considered acidic, they are slightly basic and ranged from 7.58 to 7.87 and are in the acceptable range for salmonid and macroinvertebrate species. Water temperature and DO are two of the most critical factors in determining salmonid survival. For many species of salmonids, exposure to low levels of DO (less than 5.0 - 6.0 mg/l) and exposure to high water temperatures (above 25°C) can result in mortality. Water temperature and DO were within the optimal range for spawning, egg development and growth. In situ water quality measurements of DO ranged from 10.06 mg/l to 12.90 mg/l and water temperature ranged from 5.37 °C to 10.27°C, indicating a cold water and oxygen-rich environment.

Table 1. In Situ Water Quality Measurements

Station	Stream Name	Date	T (°C)	pН	DO (mg/L)	DO (% sat.)	Conductivity (µmhos/cm)
BDC-01	Big Deer Creek	9/4/12	8.36	7.58	11.45	97.3	108
BDC-02	Big Deer Creek	9/4/12	10.27	7.87	10.24	95.9	84
UBC-01	Upper Blackbird Creek	9/5/12	5.37	7.77	12.90	102,6	75
FWFP	Freshwater Fish Pond	9/5/12	8.21	7.72	10.06	86.0	75

4.2 MACROINVERTEBRATES

Overall, 65 taxa were collected from 3 samples (Appendix E). Macroinvertebrates were identified to the lowest practical taxon. As discussed previously, the choice of core metrics was consistent with the Stream Macroinvertebrate Index (SMI) developed by the IDEQ to determine appropriate life use support in Idaho's streams.

The results of the macroinvertebrate collection in riffle habitat show that:

- The number of all macroinvertebrates was 228, 447 and 500, at BDC 01 T1, T2 and T3, respectively;
- Idaho SMI basin ratings were rated "very good" for all three macroinvertebrate transects. TI was rated 94.64, T2 was rated 99.84 and T3 was rated 97.12;
- The Hilsenhoff Biotic Index (HBI) derives a community-based estimate of overall pollution at a given site. HBI tolerance values range from 0 -10, with 0 being the most pollution intolerant and 10 being the most pollution tolerant taxa. HBI values were 2.81, 3.19 and 3.13 for T1, T2 and T3 respectively. The low HBI values indicate the presence of sensitive (intolerant) organisms;
- The Metals Tolerance Index (MTI) is based upon a correlation of invertebrate species present in known metals contaminated streams versus those present in unimpacted streams. MTI values were 2.57, 2.58 and 2.51 for T1, T2 and T3 respectively. The low MTI values indicate the presence of sensitive (intolerant) organisms to metals contamination;
- The Fine Sediment Biotic Index (FSBI) is a measure of the number of species present that are
 tolerant of increased sediment in the stream substrate. FSBI values were 120, 153 and 144 for
 T1, T2 and T3. The high FSBI indicates there are more sediment tolerant species present in the
 sample;
- The Shannon-Weaver index is a measure of the number of species (i.e., diversity) and the number of individuals within each species (i.e., evenness). The Shannon-Weaver index values were 4.69, 4.79 and 4.42 for T1, T2 and T3. A Low Shannon-Weaver index value indicates low species diversity;
- The EPT Index, a summation of taxa in the pollution-sensitive Ephemeroptera (mayflies),

Plecoptera (stoneflies), and Trichoptera (caddisflies), is a richness measure specifically focusing on the presence/absence of pollution-sensitive fauna. The EPT values were 56.14, 56.60 and 48.00 percent for T1, T2 and T3 respectively. Mayflies, stoneflies and caddisflies are associated with healthy streams, the percent of mayflies, caddisflies and stoneflies indicate that water pollution is currently not a problem;

- A composition measure, percent Chironomidae plus Oligochaeta represents the numerical abundance of pollution-tolerant midges and aquatic worms. In a healthy, balanced macroinvertebrate community, percentages of pollution-tolerant organisms are minimal. This was the case in BDC01, percent Chironomidae plus Oligochaeta were 10.53, 19.91 and 10.20 at T1, T2 and T3 respectively;
- All identifiable macroinvertebrates were dried and weighted to calculate dry biomass. Extremely
 low biomasses were encountered at each macroinvertebrate transect. Biomass was 0.055 grams,
 0.157 grams and 0.118 grams for T1, T2 and T3 respectively.

4.3 HABITAT ASSESSMENT SUMMARY

Numeric habitat ratings were developed for BDC - 01 and BDC - 02 using IDEQ BURP Habitat Assessment Protocol. Using this method, seven qualitative instream variables were evaluated for the entire sampling reach. Instream habitat parameters were each scored separately and individual habitat scores were summed to provide a total habitat score. Instream habitat scores present current habitat conditions of the stream, without considering historical or future conditions. Physical habitat scores for BDC - 01 and BDC - 02 are presented in Table 2. Additional instream characterization was conducted to assess potential fish and macroinvertebrate habitat. Additional characterization scores are not summed, are not added to the overall instream habitat scores and are identified in Table 2 with an asterisk.

The following habitat conditions were noted:

- Total habitat scores were 81 and 78 (out of a possible 115) for BDC 01 and BDC 02 respectively;
- The overall instream physical habitat conditions were considered very good for the two reaches;
- The instream conditions were similar for both Big Deer Creek reaches, including a substrate dominated by cobble with riffle embeddedness ranging between 25 to 50 percent;
- In both reaches pool variability was excellent, consisting of an even mix of deep and shallow pools;

- Pool substrate in both reaches was comprised of cobble and gravel with a soft clay or mud bottom;
- Large woody debris which provided stream stability and fish habitat was qualitatively assessed and was medium at BDC 02 and high at BDC 01;
- Instream cover for fish was optimal in both reaches. In BDC 01 instream cover was greater than 50 percent, while instream cover in BDC 02 was 40 percent;
- BDC 01 was absent of overhanging canopy cover which reduces allochtonous input and can increase stream temperature. BDC 02 had a good mix of overhanging canopy.

Table 2. Physical Habitat Assessment Data

Table 2.	Physical Habita	t Assessi	ment Da	ita		
NI NI		BDC	BDC	T1	T2	Т3
Habitat Parameter		-01	-02			
Instream cover (for fish)		18	17			
Embeddedness (in riffles)		11	10			
Channel Shape		3	3			
Disruptive Pressures (on streamb	anks)	9	9			
Zone of Influence (width of ripari	an zone)	8	8			
Pool Substrate Characteristics		15	15			
Pool Variability		17	16			
Canopy Closure - BDC -01 *	Left Bank			03	02	14
	Center Up			00	00	00
	Center Down			00	00	00
	Right Bank			02	00	00
Canopy Closure - BDC-02 *	Left Bank			05	05	08
	Center Up			06	08	05
	Center Down			06	05	06
	Right Bank			09	05	04
Pool Count *		11	8			
Large Organic Debris (LOD) * 1		33	11			
Rosgen Stream Type *		В	В			
Submerged Cover (%) *		40	35			
Undercut Banks (%) *		15	10			
Average Pool Depth (ft) * 2		2.2	2.9			
Overhead Cover (%) *		15	45			
Stream Sinuosity *		M	M			
Total Habitat Score		81	78		_	

^{*} Score does not count towards habitat assessment

4.4 STREAM DISCHARGE SUMMARY

¹LOD has a diameter greater than 10 centimeters (4 inches) and a length greater than one meter (39 inches)

M -Riffle-dominated, moderate sinuosity

² Average pool depth is based on four representative pool measurements

Using a Marsh-McBirney Digital Flo-Mate 2000 Digital Flow Meter, twenty stream discharge measurements were collected downstream of transect 3 in BDC - 01. Stream discharge measurements are provided in Table 3. Stream discharge was measured during summer base flow which is a measure of minimum stream size and an indicator of potential fish and macroinvertebrate habitat. Total stream discharge was 7.88 cubic feet per second (CFS) which is optimal for fish spawning and fish migration paths. Flow patterns also showed a positive relationship between the proportion and abundance of stoneflies, mayflies and caddisflies collected. This is not surprising, as these species are known in general to require cold, well-oxygenated, flowing water (Merritt et al., 2008; Wiggins, 1996; Stewart & Stark, 2002).

Table 3. Stream Discharge Measurements – BDC - 01								
Tape (ft)	Width (ft)	Depth (ft)	Area (ft²)	Velocity (ft/sec)	Total Discharge (ft/sec)			
(LWE) 0	:							
1.3	0.55	0.04	0.02	0.3	0.01			
2.0	0.65	0.13	0.08	0.94	0.08			
2.7	0.50	0.46	0.23	1.22	0.28			
3.4	0.45	0.43	0.19	1.67	0.32			
4.1	0.40	0.69	0.28	1.96	0.55			
4.8	0.45	0.78	0.35	1.91	0.67			
5.5	0.50	0.69	0.35	1.18	0.41			
6.2	0.50	0.72	0.36	2.06	0.74			
6.9	0.50	0.75	0.38	1.91	0.73			
7.6	0.50	0.66	0.33	1.82	0.60			
8.3	0.50	0.56	0.28	1.80	0.50			
9.0	0.45	0.59	0.27	1.50	0.41			
9.7	0.40	0.56	0.22	1.69	0.37			
10.4	0.45	0.59	0.27	1.55	0.42			
11.1	0.50	0.52	0.26	1.52	0.40			
11.5	0.50	0.56	0.28	1.37	0.38			
12.0	0.50	0.59	0.30	1.18	0.35			
12.7	0.50	0.56	0.28	1.32	0.37			
13.5	0.50	0.43	0.22	0.89	0.20			
14.7	0.65	0.20	0.13	0.72	0.09			
(RWE) 0								
	To	tal Discharge (CFS)		7.88			

4.5 SUBSTRATE SIZE CHARACTERIZATION

Stream substrate is the site of most biotic activity such as algae growth, insect growth and development,

fish egg incubation, and small fish refuge (Davis et al., 2001). Fine sediment and its accumulation can be detrimental to salmonid spawning (a beneficial use) since it may limit the quality and quantity of the inter-gravel spaces that are critical for egg incubation (Maret et al., 1993; Scrivener et al., 1989; Young et al., 1991). Following the sample design and analysis employed by IDEQ BURP, a modified Wolman Pebble Count (Wolman et al., 1954) was conducted to quantify substrate size and composition at the same three riffle transects where macroinvertebrate samples were collected. At each transect, substrate particles (e.g., cobble, sand, gravel, etc.) were selected at evenly spaced intervals across each transect (left, left middle, middle, right middle, and right), were measured (to the nearest millimeter [mm]), and were recorded on Wolman Pebble Count Data Sheets (Appendix F). A minimum of 50 particle measurements were made per riffle.

Wolman Pebble Count Data revealed predominate substrate size consisted of Very Coarse Pebbles (31.1 mm to 64 mm), Small Cobble (64.1 mm to 128 mm), and Large Cobble (128.1 mm to 256 mm). The mean substrate size was calculated and ranged between 74.4 mm to 149.3 mm. Cobble-sized substrate provides the greatest amount of usable habitat to benthic macroinvertebrates. However, the embeddedness (the degree to which very coarse pebbles and larger sizes of particles (cobbles and boulders) are surrounded or covered by fine sediment in riffle and pool habitat ranged from 20 to 90 percent and embeddedness averaged 55 percent.

4.6 FISH TISSUE CONCENTRATIONS

In accordance with the IDEQ mercury water quality criteria guidance (IDEQ, 2005), all fish tissue samples were analyzed as fillets without skin. Each individual fillet sample was homogenized and was analyzed for total mercury, aluminum, arsenic, cadmium, cobalt, lead, manganese, nickel selenium, thallium and zinc. Fish tissue samples were first analyzed for total mercury and were to be only analyzed for methylmercury if total mercury concentrations were within 20 percent of the IDEQ fish tissue mercury criterion of 0.3 milligrams per kilogram on a wet weight basis (mg/kg ww). A summary of analytical results for each subset of metal compounds is provided below. A copy of the fish tissue analytical report is provided on compact disk.

Mercury

EPA Method 1631, Revision E, Mercury in Water by Oxidation, Purge and Trap, and Cold Vapor

Atomic Fluorescence Spectrometry was used for the determination of mercury (dry & wet weight) in fish tissue. Mercury was detected in all fish tissue samples. Dry weight mercury concentrations in BDC - 01 ranged from 0.0774 milligrams per kilogram (mg/Kg) to 0.300 mg/kg. The average dry weight mercury concentration was 0.185 mg/Kg. Dry weight mercury concentrations in BDC - 02 ranged from 0.166 mg/Kg to 0.498 mg/Kg. The average dry weight mercury concentration was 0.296 mg/Kg. Dry weight mercury concentrations in UBC - 01 ranged from 0.154 mg/Kg to 0.618 mg/Kg. The average dry weight mercury concentration was 0.258 mg/Kg.

Wet weight mercury concentrations in BDC - 01 ranged from 0.01 mg/Kg to 0.06 mg/kg. The average wet weight mercury concentration was 0.04 mg/Kg. Wet weight mercury concentrations in BDC - 02 ranged from 0.03 mg/Kg to 0.103 mg/Kg. The average wet weight mercury concentration was 0.06 mg/Kg. Wet weight mercury concentrations in UBC - 01 ranged from 0.03 mg/Kg to 0.117 mg/Kg. The average wet weight mercury concentration was 0.05 mg/Kg. Mercury concentrations in all fish tissue samples were below the IDEQ fish tissue mercury criterion of 0.3 mg/Kg on a wet weight basis (mg/Kg ww). Therefore, methylmercury concentrations in fish tissue were not analyzed. A complete summary of all metal results are presented in Tables 4 through 13 (Appendix G).

Selenium

EPA Method 7742, Atomic Absorption, Borohydride Reduction was used for the determination of selenium (dry & wet weight) in fish tissue. Selenium was detected in all fish tissue samples. Dry weight selenium concentrations in BDC – 01 ranged from 0.63 mg/Kg to 2.08 mg/Kg. The average dry weight selenium concentration was 1.58 mg/Kg. Dry weight selenium concentrations in BDC - 02 ranged from 0.29 mg/Kg to 2.66 mg/Kg. The average dry weight selenium concentration was 1.67 mg/Kg. Dry weight selenium concentrations in UBC - 01 ranged from 0.51 mg/Kg to 1.93 mg/Kg. The average dry weight selenium concentration was 1.42 mg/Kg. Wet weight selenium concentrations in BDC – 01 ranged from 0.14 mg/Kg to 0.48 mg/Kg. The average wet weight selenium concentration was 0.347 mg/Kg. Wet weight selenium concentration was 0.424 mg/Kg. Wet weight selenium concentrations in UBC - 01 ranged from 0.12 mg/Kg to 0.40 mg/Kg. The average wet weight selenium concentration was 0.292 mg/Kg. A complete summary of all metal results are presented in Tables 4 through 13 (Appendix G).

Trace Metals

EPA Method 200.8, Inductively Coupled Plasma – Mass Spectrometry was used for the determination of trace metals (dry & wet weight) in fish tissue. For the Idaho Cobalt project, trace metals consist of aluminum, arsenic, cadmium, cobalt, lead, manganese, nickel, thallium and zinc. Trace metals were detected in extremely low concentration ranges in all fish tissue samples. A complete summary of all metal results are presented in Tables 4 through 13 (Appendix G).

4.7 CONCLUSIONS

In situ water quality measurements indicate pH values are not acidic, they are slightly basic and ranged from 7.58 to 7.87 and are in the acceptable range for fish and macroinvertebrate species. Water temperature and DO, two of the most critical limiting factors for fish and macroinvertebartes are within the optimal range for spawning, egg development and growth. In situ water quality measurements of DO ranged from 10.06 mg/l to 12.90 mg/l and water temperature ranged from 5.37 °C to 10.27°C, indicating a cold water and oxygen-rich environment.

Overall, 65 macroinvertebrate taxa were collected from 3 samples in BCD – 01. The Idaho SMI basin ratings were rated "very good" for all three macroinvertebrate samples, T1 was rated 94.64, T2 was rated 99.84 and T3 was rated 97.12. Of the individual core metrics, percent EPT revealed there is a good presence of pollution-sensitive fauna indicating good water quality. Conversely, the percent Chironomidae plus Oligochaete were low indicating the abundance of pollution-tolerant midges was low. Other metrics sensitive to pollution and metals contamination, including HBI and MTI were also low indicating water pollution and metals contamination was low. The Fine Sediment Biotic Index (FSBI) was high, ranging from 120 to 153, indicating there are more sediment tolerant species present. The Shannon-Weaver index revealed low species diversity. In addition to low species diversity, macroinvertebrate biomass was extremely low ranging from 0.055 grams to 0.118 grams.

Total habitat scores were 81 and 78 (out of a possible 115) for BDC - 01 and BDC - 02 respectively. The instream conditions were similar for both Big Deer Creek reaches, including a substrate dominated by cobble with riffle embeddedness ranging between 25 to 50 percent. In both reaches pool variability was excellent, consisting of an even mix of deep and shallow pools. Instream cover for fish was optimal in both reaches. In BDC – 01 instream cover was greater than 50 percent, while instream cover in BDC – 02 was 40 percent.

Thirty individual fish tissue samples were analyzed for total mercury, aluminum, arsenic, cadmium,

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

cobalt, lead, manganese, nickel selenium, thallium and zinc. Low concentrations of metals including mercury were detected in all fish tissue samples. Mercury concentrations were below the IDEQ fish tissue mercury criterion of 0.3 milligrams per kilogram on a wet weight basis (mg/kg ww) and therefore methylmercury concentrations were not analyzed.

5.0 REFERENCES

Davis, J., G. Minshall, C. Robinson and P. Landres. 2001. Monitoring Wilderness Stream Ecosystems. Report General Technical Report, RMRS-GTR-70. U.S. Dept. of Agriculture, U.S. Forest Service, Roc ky Mountain Research Station. Ogden, UT. 137 pp.

Dieterich, M. 1992. Insect community composition and physico-chemical processes in summerdry streams of western Oregon. Ph. D. thesis, Oregon State University, Corvallis OR, 191 pp.

Environ International Corp. (ENVIRON). 2009. Biological Assessment and Monitoring Program. Revision 2 November 2009

Feminella, J. W., 1996. Comparison of benthic macroinvertebrates assemblages in small streams along a gradient of flow permanence. Journal of the North American Benthological Society 15: 651–669.

Green, J., M. Passmore, and H. Childers. 2000. A Survey of the condition of streams in the primary Region of Mountaintop Mining/Valley Fill Coal Mining. U.S. EPA, Region 3, ESD, Aquatic Biology Group. Wheeling, West Virginia.

Idaho Department of Environmental Quality (IDEQ). 2005. Implementation Guidance for the Ida ho Mercury Water Quality Criteria. Idaho Department of Environmental Quality. Boise, ID, April.

Idaho Department of Environmental Quality (IDEQ). 2007. Beneficial Use Reconnaissance Program Field Manual for Streams. Boise, Idaho.

Kaufm a nn, P.R. and E.G. Robison. 1998. Physical Habitat Characterization. pp 77-118 In: J.M. Lazorchak, D.J. Klem m and D.V. Peck (eds.)). Environmental Monitoring and assessment Program — Surface W a ters: Field Operations and Methods Manual for Measuring the Ecological Condition of W a deable Streams. U. S. Environmental Protection Agency, Office of Research and Development, Wa shington, D.C. EPA/620/R-94/004F.

Kaufm a nn, P.R., P. Levine, E.G. Robison, C. Seeliger, and D.V. Peck. 1999. Quantifying Physical Habitat in Wa deable Streams. U.S. Environmental Protection Agency, W a shington, D.C. EPA/620/R-99/003.

Maret, T., T. Burton, G. Harvey and W. Clark. 1993. Field Testing of New Monitoring Protocols to Assess Brown Trout Spawning Habitat in an Idaho Stream. North American Journal of Fisheries Management 13(3): 567-580.

Merritt, R. W., Cummins, K. W., and Berg, M. B. (eds.). 2008. An introduction to the aquatic insects of North America, 4thed. Kendall/Hunt Publishing Company, Dubuque IA, 1158 pp.

Minshall, G.W., K.W. Cummins, RC Petersen, C.E. Cushing, D.A. Bruns, J.R. Medell, and R.L. Vannote. 1985. Developments in stream ecosystem theory. Can. J. Fish. Aquat. Sci. 42: 1045-1055.

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

ohnson, R., T. Wiederholm and D. Rosenberg. 1993. Freshwater biomonitoring using individual organisms, populations, and species assemb lages of benthic macroinvertebrates. In Freshwater biomonitoring and benthic macroinvertebrates. V. H. R. D.M Rosenberg (ed.). Chapman and Hall. New York, NY. 4. 40-158.

Stewart, K. W. and Stark, B. P. 2002. Nymphs of North American stonefly genera (Plecoptera). The Caddis Press, Columbus OH. 510 pp.

Telesto Solutions, Inc. (Telesto). 2010. NPDES Quality Assurance Project Plan for the Idaho Cobalt ProjectWiggins, G. B. 1996. Larvae of the North American caddisfly genera (Trichoptera), 2nd ed. University of Toronto Press, Toronto Canada. 457 pp.

Wolman, M.G. 1954. A Method of Sampling Coarse River-bed Material. Transactions of American Geophysical Union 35: 951-956.

APPENDIX A

YSI CALIBRATION RESULTS



INSTRUMENT CALIBRATION REPORT

Pine Environmental Services, Inc.

7332 S. Alton Way, Bldg. 13, Suite E. Centennial, CO 80112 Toll-free: (866) 960-PINE (7463)

Pine Environmental Services, Inc.

Instrument ID 18971
Description YSI 556
Calibrated 8/28/2012

Manufacturer YSI
Model Number 556
Serial Number/ Lot 11J102923
Number
Location Colorado

Department

Test Performed: Yes

Status Pass Temp °C 24 Humidity % 25

As Left Result: Pass

State Certified

<u> </u>							
		<u>Calib</u>	ration Specific	<u>ati</u> on <u>s</u>			
Gro	up# 1			Range Acc %			
Group N	ame PH			Reading Acc %	3.0000		
Stated A	Accy Pct of Re	ading		Plus/Minus	0.00		
Nom In Val / In Val	In Type	Out Val	Out Type	Fnd As	Lft As	<u>Dev%</u>	Pass/Fail
7.00 / 7.00	PH	7.00	PH	7.00	7.00	0.00%	Pass
4.00 / 4.00	PH	4.00	PH	4.00	4.00	0.00%	Pass
10.00 / 10.00	PH	10.00	PH	10.00	10.00	0.00%	Pass
Gro	ար# 2			Range Acc %	0.0000		
	lame Conducti	vity		Reading Acc %			
•	Accy Pct of Re	•		Plus/Minus	0.000		
Nom In Val / In Val	In Type	Out Val	Out Type	Fnd As	Lft As	Dev%	Pass/Fail
	ms/cm	1.413	ms/cm	1.413	1.413	0.00%	Pass
Gro	oup# 3			Range Acc %	0.0000		
	Name Redox (C	ORP)		Reading Acc %	3.0000		
Stated	Accy Pct of Re	eading	1	Plus/Minus	0.00		
Nom In Val / In Val	In Type	Out Val	Out Type	Fnd As	Lft As	Dev%	Pass/Fail
	mv	240.00	mv	240.00	240.00	0.00%	Pass
Gre	oup# 4	***************************************		Range Acc %	0.0000		
	Name Disolved	Oxygen Span		Reading Acc %	3.0000		
Stated Accy Pct of Reading				Plus/Minus	0.00		
Nom In Val / In Val	In Type	Out Val	Out Type	Fnd As	Lft As	Dev%	Pass/Fail
		100.00	%	100.00	100.00	0.00%	Pass
Gre	oup # 5						
l .	Name Dissolve	d Oxygen Zero					

As Found Result: Pass

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

APPENDIX B

SCIENTIFIC COLLECTION PERMIT AND REPORTING DOCUMENTS



Idaho Dept. of Fish and Game 2012 Scientific Collecting Permit Permit # F-12-12-12 - Greg Hildebrand, Hildebrand & Associates, LLC

ISSUED TO: Greg Hildebrand

c/o Hildebrand & Associates, LLC

4623 E Pegasus Ct Boise, ID 83716 Start Date: 07/02/2012

Expiration: 12/31/2012 Doc# 749-12-000057

You and your sub-permittees are authorized to sample and/or collect fish in the below listed drainages utilizing the prescribed methods identified in this permit. You must be present during all collection activities. This permit must be in your or your sub-permittee's possession while in the field or transporting biological samples.

SUB-PERMITTEES: Amy Chadwick

PURPOSE: Specimen collection, presence/absence, density estimate, vouchering (up to stated limits)

SPECIES: Rainbow trout; brook trout; cutthroat trout; brown trout; up to thirty (30) total may be vouchered

APPROVED METHODS OF SAMPLING: Electrofishing, seining, hook-and-line

GEOGRAPHIC AREAS OR WATERS: Salmon National Forest - Big Deer Creek, South Fork Big Deer Creek, Blackbird Creek

DISPOSITION OF SAMPLES: All live nontarget fish will be released back into the water where they were captured. Whole fish samples will be shipped to Columbia Analytical Services, Inc. in Kelso, WA. See Condition #2 below concerning bull trout mortalities.

PERMIT PROVISIONS:

- This permit is not transferable, nor may its authority be delegated. It shall be produced for inspection
 upon request of any conservation officer or other authorized representative of the Idaho Department of
 Fish and Game. Any abuse or misuse of privileges granted by this permit shall be grounds for
 revocation.
- YOU ARE WORKING IN WATERS WHERE BULL TROUT ARE KNOWN TO EXIST: this permit authorizes you as an agent of Idaho Department of Fish and Game to "take" bull trout. Take is defined as: observe, harass, capture, handle/tag, mark, measure, release, and indirect mortality. Any observations or handling of bull trout must specifically be reported in your permit report. Bull trout mortalities will be reported to Scott Grunder (scott.grunder@idfg.idaho.gov) at IDFG within 72 hours. The report will include location, size, and mortality factor. As per USFWS direction, bull trout mortality may be disposed of in the stream where captured.
- No electrofishing is allowed in known bull trout spawning areas between August 15 and September 15.
 If spawning activity is observed during sampling, please document the location and avoid the area.
 Please note this information on your end-of-year report.
- 4. <u>Upon completion of work or upon expiration of this permit and/or prior to any request for renewal, a report shall be submitted within 30 days, to Idaho Department of Fish and Game, Bureau of Fisheries, 600 S. Walnut, P.O. Box 25, Boise, ID 83707, providing useful and comparable information on fish collected or waters sampled. Forms are included to assist in providing information.</u>

Questions: Fisheries Bureau - 208-334-3791

Or david.parrish@idfg.idaho.gov

Idaho Dept. of Fish and Game 2012 Scientific Collecting Permit

Permit # F-12-12-12 - Greg Hildebrand, Hildebrand & Associates, LLC

- All stationary equipment used to collect fish and wildlife (nets, traps, etc.) will have an attached metal 5. tag bearing, in legible English, the name and current address of the permit holder.
- NO COLLECTIONS SHALL BE MADE UNDER THIS PERMIT UNTIL THE LOCAL CONSERVATION 6. OFFICER OR THE PANHANDLE REGIONAL OFFICE (TOM CURET) IS NOTIFIED WHERE AND WHEN THE COLLECTION IS TO BE MADE. THIS MUST BE DONE WITHIN 48 HOURS OF SAMPLING. A record of dates, times and persons notified shall be kept and submitted at the end of the year as part of the collecting report.
- A valid fishing license is needed to conduct angling surveys. 7.

Cc: Salmon Regional Office 208-756-2271 208-769-6274 (fax)

Virgil Moore, Director

Vig Q moore

June 29, 2012

Date Issued

Questions: Fisheries Bureau - 208-334-3791

Or david.parrish@idfg.idaho.gov

Permit Number: F-12-12-12									
Permit Holder: Greg Hildebrand Affiliation: Hildebrand & Associates, LLC									
A standard inventory data set will be required for any fish community or presence/absence sampling survey within the state. Additional data is welcomed; however, this minimum amount of information must be collected for each transect (site), and the form(s) returned as part of the required report. DO NOT COMBINE TRANSECTS ONTO ONE FORM EVEN IF IN THE SAME STREAM. If your report/data is compatible with the requirements of this form, they will be accepted. Please write legibly. FAILURE TO COMPLETE THIS FORM AS REQUIRED MAY BE GROUNDS FOR FUTURE PERMIT DENIAL.									
Stream Name: Big Deer Creek Transect (Sample Site) No.: BDC 01 Collection Date: 09-04-12									
Location: Big Deer Creek, SAlmon-Challis NF - Downstream of Proposed Discharge Outfall (DO NOT LEAVE BLANK) (GPS Coordinates of lower end of sampling reach − Latitude, Longitude, or UTM with zone datum) 45.10'0.1.1" 114.21'47.1" Estimator (√): Direct observation (Snorkeling) Mark/Recapture Presence/Absence Depletion (one or more electrofish passes) Flook & Line Water Temp. (C°): 8.36 Water Temp. (C°): 8.36									
Transect Length (M):137									
Optional information on permanent fish barriers: Location: N/A Height: N/A if manmade – type: N/A									
NOTICE OF COLLECTION:									
IDFG Person Contacted:Tom Curet Date & Time of Contact: 8-31-12: 14:26									
Disposition of Fish: Columbia Analytical Services									
Report densities/CPUE in the table below. Indicate type of gear used. Record fishing effort and age class by species.									
Fish Species Gear Type Used Time Fished #fish/100m² or Age Class (specify column) CPUE No. of 0 year No. of ≥ 1 year									
Rainbow Trout Hook & Line 1 Hour 10 Fish/Hr Zero 10									
Rambow Hour House Enter									

NUMBERS OF NON-GAME FISH OBSERVATIONS SHOULD BE RECORDED ON ELECTROFISHING DATA SHEET

Stream Name: Big Deer Creek	_ Transect (Sample Site) No.: BDC 01	Collection Date: <u>09-04-12</u>
Location: Inside Salmon-Challis NF	Crew Leader: <u>Greg Hildebrand</u>	Permit No.: F-12-12-12

Length Range	Sı	pecies			Spec	cies	
(mm)	Rainbow		(mm)				
/			310-319				
			320-329				
			330-339				
50-59			340-349				
60-69			350-359				
70-79			360-369				
80-89			370-379				
90-99			380-389				
100-109			390-399				
110-119			400-409				
120-129			410-419		<u>_</u>		
130-139			420-429				
140-149			430-439				
150-159			440-449				
160-169			450-459				
170-179			460-469				
180-189	II		470-479				1
190-199	11		480-489				
200-209	II		490-499				×
210-219	ī		500-509				i
220-229	II		510-519				1
230-239			520-529				
240-249			530-539				
250-259			540-549				1
260-269							
270-279							
280-289			Total	10			
290-299			Each	Rainbow			
300-309			Species	Trout			
200-208				1	aga tabla		

Density estimates are to be calculated and entered on the front page table.

Other species sampled/observed:_	N/A					<u> </u>	
----------------------------------	-----	--	--	--	--	----------	--

Permit Number: F-	12-12-12								
Permit Holder: Gre	eg Hildebrand	Affil	iation: Hildebrand &	Associates, LLC					
within the state. A for each transect TRANSECTS ON with the requirement	dditional data is we (site), and the for NTO ONE FORM ents of this form, the	elcomed; however, m(s) returned as EVEN IF IN TH ev will be accepte	fish community or this minimum amore part of the require E SAME STREAN d. Please write legi FOR FUTURE PE	ed report. DO N 1. If your report/d 1. Is your report/d	OT COMBINE ata is compatible				
Stream Name: Big D	Deer Creek	Transect (San	nple Site) No.: BDC 02	2 Collection I	Date: 09-04-12				
 _		OO NOT LE	f Proposed Discharge AVE BLANK)						
(GPS Co	ordinates of lower en	d of sampling reach	- Latitude, Longitude	e, or UTM with zone	e datum)				
Estimator (1): Direct	t observation (Snork ore electrofish passe	eling) 🔲 Mark/l	Recapture Pres	ence/Absence 🗌					
Transect Length (M (At least one pool/ri): 100 ffle/run complex.)	Transact Width (l (At least 4 measure	M): 2.5 ements for an average.)					
Optional informatio Location: N	on on permanent fish	barriers: _ Height: <u>N/A</u>	if manmade -	type: N/A					
NOTICE OF COLL	ECTION:								
IDFG Person Conta	icted: Tom Curet	[Date & Time of Conta	ct: <u>8-31-12: 14:26</u>					
Disposition of Fish:	Columbia Analytica	al Services							
Report densities/CF	PUE in the table below	w. Indicate type of	gear used. Record fisl	ning effort and age cl	ass by species.				
Fish Species	Gear Type Used	Time Fished	#fish/100m² or CPUE	Age Class (sp No. of 0 year	No. of ≥ 1 year				
Rainbow Trout	Hook & Line	80 Minutes	10 Fish/Hr	Zero	10				
	,								
			The Condens Of the Sec	TROPICURIC DATA C	JCGT***				
***NUMBERS OF N	ON-GAME FISH OBSER	EVATIONS SHOULD B	E RECORDED ON ELEC	INOTISHING DATA SI					

¹

Stream Name:	Big Deer Creek	Transect (Sample Site) No.:	BDC 01	Collection Date: <u>09-04-12</u>	_

Location: Inside Salmon-Challis NF	_ Crew Leader:_	Greg Hildebrand	Permit No.: F-12-12-12
------------------------------------	-----------------	-----------------	------------------------

Length Range		Species	Species				
(mm)	Rainbow		(mm)				
			310-319				<
			320-329				1/4
			330-339				
50-59			340-349				
60-69			350-359				
70-79			360-369				
80-89			370-379				
90-99			380-389				
100-109			390-399				
110-119			400-409				
120-129			410-419				
130-139			420-429				
140-149			430-439				
150-159			440-449		·		
160-169			450-459				
170-179			460-469				
180-189	11		470-479				
190-199	11		480-489		*		
200-209	IIII		490-499				
210-219	II		500-509				
220-229	11		510-519				
230-239	-		520-529				
240-249			530-539				
250-259			540-549				
260-269							
270-279							
280-289			Total	10			
290-299			Each	10 Rainbow			
300-309			Species	Trout			

Density estimates are to be calculated and entered on the front page table.

0.1 1.1/-baseneds	2.27.4				
Other species sampled/observed:	N/A		 	·	-

Permit Number: F-	12-12-12									
Permit Holder: Gre	g Hildebrand	Affil	iation: Hildebrand &	z Associates, LLC						
for each transect TRANSECTS ON with the requirement	dditional data is we (site), and the for TO ONE FORM of this form, the	elcomed; however, m(s) returned as EVEN IF IN TH ey will be accepte	fish community or this minimum amor part of the require E SAME STREAN d. Please write legi FOR FUTURE PE	unt of information is ded report. DO No. 1 If your report/dibly. FAILURE TERMIT DENIAL.	OT COMBINE ata is compatible O COMPLETE					
Stream Name: <u>Upp</u>	Stream Name: Upper Blackbird Creek Transect (Sample Site) No.: UBC Collection Date: 09-05-12									
(GPS Coo	ordinates of lower en 45.07'01.50" 114.2 t observation (Snorke ore electrofish passes	(DO NOT LE.d of sampling reach 0'47.26" eling) ☐ Mark/l	 Latitude, Longitude Recapture ☐ Pres 		e datum)					
Transect Length (M) (At least one pool/ri): 100 ffle/run complex.)	•	M): 2 ements for an average	.)						
Optional information Location: N	n on permanent fish \ /A	Height: N/A	if manmade -	type: N/A						
NOTICE OF COLL	ECTION:									
IDFG Person Conta	cted: Tom Curet		Date & Time of Conta	ct: <u>8-31-12: 14:26</u>						
Disposition of Fish:	Columbia Analytica	I Services								
Report densities/CP	UE in the table below	w. Indicate type of g	gear used. Record fish	hing effort and age cl	ass by species.					
Fish Species	Gear Type Used	Time Fished	#fish/100m ² or CPUE	Age Class (sp No. of 0 year	No. of ≥ 1 year					
Rainbow Trout x	Hook & Line	2 Hours	5 Fish/Hr	Zero	10					
Cutthroat Hybrid										

^{***}NUMBERS OF NON-GAME FISH OBSERVATIONS SHOULD BE RECORDED ON ELECTROFISHING DATA SHEET***

Idaho Department of Fish and Game Scientific Collecting Permit Mandatory Report Form

Stream Name: Upper Blackbird Creek	Transect (Sample Site) No.: UBC	Collection Date: <u>09-05-12</u>
	Constitution Constitution	Permit No.: F-12-12-12
Location: Inside Salmon-Challis NF	Crew Leader: <u>Greg Hildebrand</u>	remit No.:_ <u>F-1Z-1Z-1Z</u>

Length Range		Spec	cies				Spec	cies	
(mm)	Rainbow x	Cutthroat	Hybrid		(mm)				
					310-319			-	
					320-329				
					330-339				
50-59					340-349				
60-69					350-359				
70-79					360-369				
80-89					370-379				
90-99				-	380-389				
100-109			-		390-399				
110-119					400-409				
120-129					410-419				
130-139					420-429				
140-149					430-439				
150-159					440-449				
160-169					450-459				
170-179					460-469				
180-189	I I		<u> </u>		470-479		- -		
190-199	11				480-489				
200-209	I				490-499				
210-219	IIII				500-509				
220-229	I				510-519				
230-239					520-529				
240-249					530-539				
250-259					540-549				
260-269									
270-279									
280-289					Total	10		Ì	
290-299					Each		Cutthroat		
300-309					Species	Hybrid			

Density estimates are to be calculated and entered on the front page table.

Other species sar	npled/observed:_	N/A		 	
Ottor observe per-		13/1	 -		

APPENDIX C

FISH SPECIES COLLECTION FORMS

Idaho Cobalt Project - Fish Tissue Collection Log

Collection Method: Rod and Reel Weather: Sunny

	1	C(2) Co-2		10:00		00;05		50.00		50:0°		0.00	20.00		20:00		60:00		සිට : අන		00.09	
Location Description	Downstream of Proposed Outfall		To the of Description	Downstream of Froposed Outlan		Downstream of Proposed Outtall	11 0 0	Downstream of Proposed Outfall		Downstream of Proposed Outfall		Downstream of Proposed Outfall		10.46.11	Downstream of Proposed Outrain		Downstream of Proposed Outfall		Downstream of Proposed Outfall		Downstream of Proposed Outfall	
Length (mm)	(mim)	250	7,77		182		199		190		219		225	777		204		228		207		184
Fish	Deinbow	Tront	TIOUL	Rainbow	trout	Rainbow	Trout	Rainbow	Trout	Rainbow	Trout	Rainhow	Tront	Lrout	Rainbow	Trout	Rainbow	Trout	Rainbow	Trout	Rainbow	Trout
Fish		BDC-01-01		BDC-01-02		BDC-01-03		BDC-01-04		BDC-01-05		BDC_01_06	20-10-077		BDC-01-07		BDC-01-08		BDC-01-09		BDC-01-10	
Stream	Name	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek	Die Deer	1377 BIG	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek
Date		09-04-2012		09-04-2012		09-04-2012		09-04-2012		09-04-2012		000000000000000000000000000000000000000	09-04-2012		09-04-2012		09-04-2012		09-04-2012		09-04-2012	
Sampler		Hildebrand		Chadwick		Wilkins		.1								60						

Idaho Cobalt Project - Fish Tissue Collection Log

Collection Method: Rod and Reel Weather: Sunny

	Location Description	21 0. < 3	Upstream of Proposed Outfall		Upstream of Proposed Outfall	10, 01	Upstream of Proposed Outfall	11 3, 41	Upstream of Proposed Outfall		Upstream of Proposed Outfall		Upstream of Proposed Outfall		Tractream of Proposed Outfall	Opsileani of Loposed Curan	10.161	Upstream of Proposed Outrall		Upstream of Proposed Outfall	= 0.00	Upstream of Proposed Outfall	
	Length	(mm)		210		208		203		206		181		216		4	209		190		194		182
	Fish	Species	Rainbow	Trout	11001	Rainbow	Trout	Rainbow	Trout	Rainbow	Trout	Rainbow	Trout										
S	Fish	Д		BDC-02-01		BDC-02-02		BDC-02-03		BDC-02-04		BDC-02-05		90-CU-JUB	DDC-07-00		BDC-02-07		BDC-02-08		BDC-02-09		BDC-02-10
	Stream	Name	Big Deer	Creek	Big Deer	Jest C	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek	Big Deer	Creek								
	Date		09-04-2012		09-04-2012		09-04-2012		09-04-2012		09-04-2012	1	00 04 2012	03-04-5015		09-04-2012		09-04-2012		09-04-2012		09-04-2012	
	Sampler		Hildebrand		Chadwick		Wilkins											•					

Idaho Cobalt Project - Fish Tissue Collection Log

Collection Method: Rod and Reel Weather: Sunny

Sampler	Date	Stream Name	Fish	Fish	Length	Location Description
4			E	Species	(mm)	
Hildebrand	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-01	Cutthroat X	221	Freshwater Fish Pond
Chadwick	09-05-2012	Upper Blackbird		Rainbow-		# 1
		Creek	UBC-05	Cutthroat X	199	Freshwater Fish Pond
Wilkins	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-03	Cutthroat X	215	Freshwater Fish Pond
	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-04	Cutthroat X	218	Freshwater Fish Pond
	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-05	Cutthroat X	210	Freshwater Fish Pond
	09-05-2012	Upper Blackbird	\w_	Rainbow-		
		Creek	UBC-06	Cutthroat X	192	Freshwater Fish Pond
	09-045-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-07	Cutthroat X	214	Freshwater Fish Pond
	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-08	Cutthroat X	184	Freshwater Fish Pond
	09-05-2012	Upper Blackbird		Rainbow-		
		Creek	UBC-09	Cutthroat X	207	Freshwater Fish Pond
	09-05-2012	Upper Blackbird		Rainbow-		1
		Creek	UBC-10	Cutthroat X	185	Freshwater Fish Pond

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

APPENDIX D

CHAIN OF CUSTODY FORMS

	=
	11
	C
	0
	E
	č
	į
	Ē
	i i
1	0
1	V
M	4
4	

OF CUSTODY

SR#

(CIRCLE ONE) (Total Metals: [An] (As Sb Ba Be B Ca (A) (Cin) Cr Cu Fe (Pip) Mri (Mri) Mo (Ni) K Ag Na (Se) Sr (TI) Sn V (Zn) [Hg Above circles METALS. SAMPLES WILL BE RAN FOR TOTAL MERCURY AND WILL BE REMARKS IF MEHAL MERCUKI HASTOBE SPECIAL INSTRUCTIONS/COMMENTS: EACH TISS WE SMAYILE WILL BE ANALYZED FOR THE MPRCURY CONCENTRATIONS Š. *INDICATE STATE HYDROCARBON PROCEDURE: AK CA WI NORTHWEST OTHER: AL Ba Be B Ca Cd Co Or Cu Fe Pb Mg Mn Mo Ni K Ag Na ONLY ANKYZED FLE MEHATILMERCURY FF PLAN My ty CCINTACT FIRES OF O'S MG 1317 South 13th Ave., Kelso, WA 98626 | 360.577.7222 | 800.695.7222 | 360.636 1068 (lax) Circle which metals are to be analyzed: PLRASE WITHEN 30% Dissolved Metals: Al As Sb Service Ovatile Organics by GC/MS COMPS CO Anabarzer) NUMBER OF CONTAINERS 1. Ides RAMIS . ASSOCIATES IIC TURNAROUND REQUIREMENTS 4623 E DUSTABLUSCT Standard (15 working days) INVOICE INFORMATION 15 II 方任 TISK. 48 hr. MATRIX IST. P.T. FILL. 15th FIS. AC 17 FISK Baise In 83711 Provide FA''. Results 100 JA 100 Buil 20C 01-01 01-01 P.O. # 334/4 LAB I.D. MAIL MENUES GREG ANTELNE PRAMI & GMATL OF C. DRC COMPANY NAME HILDERS RAND & ASSOCIATE HILDE BARNI 24 hr. Bill To: 1415 TIME 330 2012 DEGASAS COSKT 8/1/B カイプか 4/2 CIMIB 0/2/ C1/h/b 21/1/2 0/4/6 STATE OF THE Report Dup., MS, MSD as Molecham Routine Report: Method 101-S19 REPORT REQUIREMENTS IV. Data Validation Report Blank, Surrogate, as CLP Like Summary GREG HINAHO 4633 E G-101 (no raw data) -01-03 300-01-10 10-10-10-10-30-10-06 10-12-XX といしい required 30C-c1-07 required 3x -c1-0 V. EDD 3 =

Copyright 2012 by ALS Group

Date/Time

Signature

Date/Time

Firm

Printed Name

Signature

Date/Ti

Frinted Name

Firm ASSOCI ATPS

Sighature Date/Time ARAND HALDERSKINN) OF Prinked Name Firm ASSACE AND

bry Wildeliano 9/10/12

RELINQUISHED BY:

RECEIVED BY:

Requested Report Date

Firm

Printed Name

RECEIVED BY:

Sample Shipment contains USDA regulated soil samples (check box if applicable)

RELINQUISHED BY:

CHAIN OF CUSTODY

(CIRCLE ONE) TOTAL MERCURY AND WALL BE (Zn/Hg) WILL BO ANNIZED FOR THE WG WIVE IF METHIL MERCUKY MAS REMARKS MERCURY CONCENTRATIONS Date/Time Sn V Sn RECEIVED BY: (Se)Sr(TJ) NORTHWEST OTHER: △ RSK 175 (Methane (CO2) Pb Mg Mn Mo Ní K Ag Na Cr Cu Fe (Pb) Mg (Min) Mo(Ni) K Ag Na Printed Name Sample Shipment contains USDA regulated soil samples (check box if applicable) Signature 1613 O RAN FER SPECIAL INSTRUCTIONS/COMMENTS: EXH TYSSUE SAMPILE 15:12 "INDICATE STATE HYDROCARBON PROCEDURE: AK CA WI THES! PAGE Date/Time :008 'Ha RELINQUISHED BY: Cu Fe Above circleis METALS , SAMPLES Will be 20% OF O.3 Mg FAL METHILMERCURT Ö Ва Ве В Са (Сd)/Сd 1317 South 13th Ave., Kelso, WA 98626 | 360.577.7222 | 800.695.7222 | 360.636.1068 (lax) Be B Ca Cd Drintod Nomo Signature Circle which metals are to be analyzed: Cas Dieseld (*See below) to be ANASYZED Ва Sp Tetal Metals: (As Sb ONLY ANALIZED RE WITHEN Semivolatile Organics by GC/MS

8250 82700 82700 GC/MS

DHAP MIS

DHAP MIS

Semivolatile Organics by GC/MS Dissolved Metals: Al As RECEIVED BY: NUMBER OF CONTAINERS HildeSpanni ASSOCIATESIIC TURNAROUND REQUIREMENTS P.O. # 23-14] Standard (15 working days) INVOICE INFORMATION Requested Report Date B GMRTL: tem 48 hr. ラリ 不以 NS.H PPO. JECT MATRIX FIST. List FIST TY. Provide FAX Results BRAND & ASSOCIATES 09-10 09-10 856 856 856 いから 500 TO (2 60 802-63 802-63 802-63 LAB I.D. P Buise ID 8374 24 hr. WILLIAMAIN 1330 8 TIME DeGASMS COBALT GREGAHING GRAND Hildeline Hildeline 09/9/13 RELIMOUISHED BY: 4 0/1/0 0/1/6 5/4/6 5/2/6 II. Report Dup., MS, MSD as ウ/ケ/タ 0/4/6 9/7/3 STAN STAN 42% 1. Routine Report: Method DATE REPORT REQUIREMENTS Data Validation Report Blank, Surrogate, as CLP Like Summary TOM (ALS) Environmental 1 200 C-ERT WITHLANN) K STO 60-(no raw data) 12/6/ ADC-62-05 BNC-63-63 BDC-CB-COL BDC-03-08 2-60-80x -3-06 BK-62-07 BDC-03-03 required required BDC-09 63 EDD SAMPL PROJECT MANAGER PHONE # 208 PHOJECT NUMBER E-MAIL ADDRESS き > = ADDRESS

ALS) Environmental

CHAIN OF CUSTODY

1317 South 13th Ave., Kelsu, WA 98626 | 360.577.7222 | 800.695.7222 | 360.636.1068 (tax)

SR SR S.7222 | 360.636.1068 (fax) PAGE | OF L

(CIRCLE ONE) HAS 72 52 (Se) Sr(TI) Sn v (Zn) Hg REMARKS HEVE CIRCLED METHS . SAMPLES WILL BE RAIN FIR TATAL MERCURY AND WILL be ONLY TISSUE SAMPLE WILL BE ANALYZED FER THE TETAL MERCURY CONCENTRATIONS RECEIVED BY: IF MeHHILMERCURY *INDICATE STATE HYDROCARBON PROCEDURE: AK CA WI NORTHWEST OTHER: A Cr Cu Fe (Pb/ Mg (Mn/ Mo(Ni) K Ag Na K Ag Na Sample Shipment contains USDA regulated soil samples (check box if applicable) \$ 5. E. E. E. 1008 14d E. RELINQUISHED BY: O Č ANTHEN JOB OF O'S MY/ ANTHEN ACON CUNTRIT FIRST FER METHYLMERCURY 000 Cq Cq SPECIAL INSTRUCTIONS/COMMENTS. EAU Pesticides/Herbi В Ca Ω Be Be Circle which metals are to be analyzed: Sas () Siesei () Sie Ва Ва Total Metals: (Al (As) Sb Semivolatile Organics by GO/MS

Semivolatile Organics by GO/MS As Dissolved Metals: Al ANALYZeis RECEIVED BY NUMBER OF CONTAINERS DEGASASCT TURNAROUND REQUIREMENTS HIDDESPEND - ASSOCIATES ILC BOAND & GMAILILLIN Standard (15 working days) INVOICE INFORMATION FIS! FIST THE THE EST. 西山 TST. TY. Requested Report Date 182-05 [IS] WAS 10 FEST P. ASSOCIATES Provide FAX Results WBC-43 MBC-64 MBC of 10-XM Bill To: 4633 巨 UBC-08 MBC-09 UBC-CA) LABI.D. MBC-CI PO.# 2344 HILLINGSRAND 537 24 hr. 5 day 0.0 Bise 3 TIME DEGABUS 3 9/S/B 1HB 9/S/B 9/2/19 9/5/B <u>₹</u> ILDEARAND 9/2/B 9/13 01/5/6 0 RELINQUISHED BY: Report Dup., MS, MSD as Routine Report: Method REPORT REQUIREMENTS IV. Data Validation Report Blank, Surrogate, as III. CLP Like Summary HJAHO STO GREGA G P. C. (no raw data) 1/36-08 1/3c-03 0,0 13c-28 60-WBC - 0 Y ري 50 required required 1 7 EDD PROJECT NUMBER -MAIL ADDRESS COMPANY NAME 4,86. NB. BC BR 0 3 <u>=</u>: > 3 ADDRESS

Printed Name Firm Copyright 2012 by ALS Group

Date/Time

Signature

Date/Time

Signature

Firm

Printed Name

Firm

Printed Name

Firm A SSGC MPRS/IC

Bate/Time

(de LOCAM)

phature

ALYSTS INC

7

		The second secon		TO AND THE PARTY OF THE PARTY O			
		EcoAnalysis Project#	社				
		Company:		Hildebrand & Associates, LLC	sociates, LLC		
		Total # of Samples this project	his project.	3 Total (T1, T2 & T3)	13)		
		# of Samples Shipped this shipment	d this shipment.	3 Total (T1, T2 & T3)	T3)		
					0		
Containers	Stream	Site #	Rep	Device Type	Dimensions	Habitat	Collection Date
100	Big Deer Ofeek	Big Deer Oreek 01	1.7	Hess Sampler	13" x 36" 500 µm	स् सिक	9/4/2012
	Bo Deer Oreak	Bro Deer Creek 01	L-1	Hess Sampler	13" x 16" 500 µm	Riffle	9/4/2012
- Cap	Sig Deer Cleek	Sig Deer Creek 01	(P)	Hess Sample:	13"× 16" 500 µm	_	9/4/2012
Relinquished Byl Date.	Mari H. Tolah	40/60 hours	913013	Condition; Scot	r. Good		
Company:	milenpland & Associates						
Received Bys Date:	- Cras	9/1	9/12 2012	Condinor	Condition: Good		
Company:	NO VICTORIAN	वादो					
Refinquished By! Date.		A CONTRACTOR OF THE PERSON NAMED IN CONT		Condition			
Company:							
Received By/ Date:				Constition		dia mindri maning tanga paga paga masabada dak di adid di didapp pipuli disersi	PREJAMENTALISTA AND VALORISTA PARTICIPATO PRESTATION PROPRIETO PROVINCIONISTA
Company:							
Relinquished By: Date:				Condition	1.		
Company:	me i inde de deste e insperimente constitución en establisha de senda de la Petrol de definida de definida establisha en establisha de deste en establisha de definida de definida establisha en establisha de deste en establisha de deste en establisha de de deste en establisha de de deste en establisha de deste en establi	annessen eigen er der erfer erfer des eigen eine de des des des des des des des des des					
Received By: Date				Condition			
Company:							

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

APPENDIX E

MACROINVERTEBRATE RESULTS

Hildebrand Idaho Cobalt ID BURP Benthos 2012 Dry Weights

ANALYSTS INC.
University of Idaho Analytical Sciences Laboratory

Units: grams Reporting Limit: 0.001

EcoA						Sample	Analysis	
Sample ID	Site ID	Rep	Collect Date	Matrix	Parameter	Weight - Dry	Method	ASL ID
6147.1-1	Big Deer Creek 01	T1	04-Sep-12	Solid - Wet Weight	Sample Weight - Dry	0.055	Gravimetric	E1202817
6147.1-2	Big Deer Creek 01	T2	04-Sep-12	Solid - Wet Weight	Sample Weight - Dry	0.157	Gravimetric	E1202818
6147.1-3	Big Deer Creek 01	T3	04-Sep-12	Solid - Wet Weight	Sample Weight - Dry	0.118	Gravimetric	E1202819

Hildebrand Idaho Cobalt ID BURP Benthos 2012 Sort Report

Eco Analysts, inc.

								Estimated	Estimated					
EcoA				Collection			Primary	Pre-Rinse	Post-Rinse			Estimated	Estimated	Estimated
Sample	D Stream	Site ID	Rep	Date	Sorter	Sort Date	Matrix	Volume (L)	Volume (L)	QC Sorter	QC Date	%Recovery1	%Recovery2	%Recovery3
6147.1-1	Big Deer Creek	Big Deer Creek 01	T1	09/04/12	Susie Patton	09/28/12	Inorganic	0.16	0.06	Megan Payne	09/28/12	100.00	N/A	N/A
6147.1-2	Big Deer Creek	Big Deer Creek 01	T2	09/04/12	Susie Patton	09/28/12	Inorganic	0.15	0.10	Megan Payne	09/28/12	100.00	N/A	N/A
6147.1-3	Big Deer Creek	Big Deer Creek 01	T3	09/04/12	Susie Patton	09/28/12	Inorganic	0.05	0.05	Megan Payne	09/28/12	100.00	N/A	N/A

Hildebrand Idaho Cobalt ID BURP Benthos 2012
Data are adjusted for subsampling
**Calculations use EcoAnalysts standard attributes

ECOANALYSTS, INC.

	Site ID Rep Collection Date	09-04-2012 Hess Sampler 100.00	Big Deer Creek Big Deer Creek 01 T2 09-04-2012 Hess Sampler 100.00 6147.1-2	Big Deer Creek Big Deer Creek 01 T3 09-04-2012 Hess Sampler 100.00 6147.1-3
Abundance Measures Corrected Abundance		228.00	447.00	500.00
EPT Abundance		128.00	253.00	240.00
Dominance Measures Dominant Taxon		Heterlimnius sp.	Zapada cinctipes	Heterlimnius sp.
Dominant Abundance		34.00	86.00	123.00
2nd Dominant Taxon 2nd Dominant Abundance		Zapada cinctipes 17.00	Heterlimnius sp. 33.00	Zapada cinctipes 41.00
3rd Dominant Taxon		Taeniopterygidae	Cinygmula sp.	Cinygmula sp.
3rd Dominant Abundance % Dominant Taxon		16.00 14.91	25.00 19.24	35.00 24.60
% 2 Dominant Taxa % 3 Dominant Taxa		22.37 29.39	26.62 32.21	32.80 39.80
Richness Measures				
Species Richness		42.00	54.00	50.00
EPT Richness Ephemeroptera Richness		20.00 7.00	29.00 11.00	26.00 11.00
Plecoptera Richness		7.00	8.00	8.00
Trichoptera Richness Chironomidae Richness		6.00 9.00	10.00 11.00	7.00 12.00
Oligochaeta Richness		0.00 33.00	0.00	0.00
Non-Chiro. Non-Olig. Richness Rhyacophila Richness		33.00 2.00	43.00 3.00	38.00 3.00
Community Composition				
% Ephemeroptera % Plecoptera		23.25 25.00	17.23 28.41	27.80 16.00
% Trichoptera		7.89	10.96	4.20
% EPT % Coleoptera		56.14 19.74	56.60 10.51	48.00 29.20
% Diptera		11.84	21.25	11.00
% Oligochaeta % Baetidae		0.00 4.39	0.00 2.68	0.00 3.60
% Brachycentridae		2.63	3.36	1.00
% Chironomidae % Ephemerellidae		10.53 6.58	19.91 5.59	10.20 11.40
% Hydropsychidae		0.44	0.22	0.40
% Odonata		0.00	0.00	0.00
% Perlidae % Pteronarcyidae		2.63 0.00	2.01 0.00	0.60 0.00
% Simuliidae		0.00	0.00	0.00
Functional Group Composition % Filterers	n	2.19	4.70	1.00
% Gatherers		32.46	34.90	45.20
% Predators % Scrapers		23.68 18.86	19.02 11.63	17.80 19.20
% Shredders		16.67	25.73	10.00
% Piercer-Herbivores % Unclassified		0.00 0.00	0.00 0.22	0.00
Filterer Richness		2.00	3.00	1.00
Gatherer Richness Predator Richness		13.00 13.00	19.00 16.00	18.00 18.00
Scraper Richness		7.00	7.00	8.00
Shredder Richness Piercer-Herbivore Richness		5.00 0.00	7.00 0.00	4.00 0.00
Unclassified		0.00	1.00	0.00
Diversity/Evenness Measures			4.44	4.22
Shannon-Weaver H' (log 10) Shannon-Weaver H' (log 2)		1.41 4.69	1.44 4.79	1.33 4.42
Shannon-Weaver H' (log e)		3.25	3.32	3.07
Margalef's Richness Pielou's J'		7.55 0.87	8.68 0.83	7.88 0.78
Simpson's Heterogeneity		0.95	0.94	0.91
Biotic Indices % Indiv. w/ HBI Value	, i	94.74	94.41	92.40
Hilsenhoff Biotic Index		2.81	3.19	3.13
% Indiv. w/ MTI Value Metals Tolerance Index		67.11	76.51	74.20 2.51
% Indiv. w/ FSBI Value		2.57 67.98	2.58 62.42	73.80
Fine Sediment Biotic Index FSBI - average		120.00 2.86	153.00	144.00 2.88
FSBI - weighted average		5.21	2.83 4.70	5.07
% Indiv. w/ TPM Value		78.95	69.80	78.00
Temp. Pref. Metric - average TPM - weighted average	-]	4.50 6.77	4.30 6.03	4.36 6.60
Karr BIBI Metrics				
Long-Lived Taxa Richness Clinger Richness	ļ	10.00 22.00	10.00 28.00	7.00 26.00
% Clingers		73.25	67.34	77.60
ntolerant Taxa Richness		19.00	26.00	22.00
	I	2 31	3 32	3.03
% Tolerant Individuals % Tolerant Taxa Coleoptera Richness		2.31 4.76 4.00	3.32 3.70 4.00	3.03 4.00 3.00

Hildebrand Idaho Cobalt ID BURP Benthos 2012
Data are NOT adjusted for subsampling
Idaho Basins SMI calculations located below TOTAL

Eco Analysts, Inc.

	Strean Site IC Rep Collection Date Device Percent Subsample EcoAnalysts Sample IC	Big Deer Creek 0 T 09-04-201; Hess Sample 1 100.00	1 Big Deer Creek 01 1 T2 2 09-04-2012 r Hess Sampler 0 100.00	Big Deer Cree Big Deer Creek 0 T. 09-04-201 Hess Sample 100.0 6147.1-
Ephemeroptera	Acentrella turbida	0	- Albara - A	1
	Baetis tricaudatus	10		17
	Caudatella hystrix	0		2
	Cinygmula sp.	4		35
	Drunella coloradensis/flavilinea Drunella doddsii	13		34
	Epeorus deceptivus/hesperus	3		2
	Epeorus grandis/permagnus	7		6
	Ephemerella sp.	0		16
	Leptophlebiidae	0 14		0
	Rhithrogena sp. Serratella tibialis	14		21 1
Plecoptera	Chloroperlidae	o o		6
stoneflies	Doroneuria sp.	6		3
	Megarcys sp.	2		1
	Paraperla sp. Perlodidae	0 2		1 4
	Sweltsa sp.	12		17
	Taeniopterygidae	16		7
	Zapada cinctipes	17		41
Coleoptera	Zapada columbiana Heterlimnius sp.	2 34		0
	Lara sp.	1	33	123
	Narpus sp.	, o		0
	Optioservus sp.	9	9	20
Diptera-Chironomidae	Zaitzevia sp.	1	4	3
nonbiting midges		0 0	5	1
	Eukiefferiella gracei gr.	3	8	1
	Eukiefferiella tirolensis	0	1	0
	Micropsectra sp.	6	20	15
	Orthocladiinae Orthocladius (Euortho.) rivulorum	0 0	0	1 0
	Orthocladius (Euorthocladius) sp.	1	0	1
	Orthocladius Complex	2	1	4
	Orthocladius sp.	3	2	2
	Potthastia gaedii gr.	4	6	11
	Rheocricotopus sp. Thienemanniella sp.	1 1	19 6	8
	Tvetenia bavarica gr.	3	20	5
	Bezzia/Palpomyia sp.	2	4	3
flies	Dicranota sp.	0	1	1
	Pericoma/Telmatoscopus sp. Rhabdomastix fascigera gr.	0	1	0
Trichoptera	Amiocentrus aspilus	Ö	1	0
caddisflies	Arctopsyche grandis	1	1	0
	Brachycentrus americanus	4	13	5
	Dolophilodes sp. Glossosoma sp.	0 4	7	0
	Lepidostoma sp.	0	5	0
	Micrasema sp.	2	1	0
	Neophylax sp.	0	0	4
	Parapsyche elsis Rhyacophila betteni gr.	0	0 1	2
	Rhyacophila brunnea gr.	6	11	2
	Rhyacophila sp.	0	6	3
	Rhyacophila vagrita gr.	1	0	0
Acari water mites	Atractides sp.	1	0	1
	Protzia sp.	4	5 1	15 2
	Sperchon sp.	2	7	9
	Stygothrombium sp.	0	1	0
	Torrenticola sp.	2	7	8
Crustacea Other Organisms		5 13	14 17	14 10
Other Organisms	TOTAL	228	447	500
				300
	IDAHO SMI BASINS SCORE IDAHO SMI BASINS RATING	94.64 Very Good *Sample 1 did not	99.84 Very Good *Sample 2 did not	97.12 Very Good
		reach the 500 count minimum for the SMI calculation		

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

APPENDIX F

WOLMAN PEBBLE COUNT

9217537565

Wolman Pebble Count

Crehetuata Data		Diffla 1	0 1			Riffle 2	2			Ri	Riffle 3	
Duositate Data	Outside Wetted	Wetted	Within	Wetted	Outside Wetted	Wetted		Within Wetted	Outside Wetted		Within Wetted	Wetted
Silt/Clay 0-1 mm	\$ 9 *	0	* *	M 0	6 E		o c	00	A23	0 (4)	17,5	0
Sand 1.1-2.5 mm	6	0	0 6	(2)	for	%	es e	Employee	**	0 5	4 m	0
Sub Total	V	Charles State of the State of t	6,000 - 2000)	9	, series .	()	60		0	on any and a second	Y
Very Fine Pebble 2.51 - 6 mm												
Pebble 615 mm											9 °	
Coarse Pebble 15.1 - 31 mm			×				N	(A)			N	and the second second
Very Coarse Pebble 31.1 - 64 mm											Z.	^-7
Small Cobble 64.1 - 128 mm			9 4 4	<u></u>			N	() ()			的效	0
Large Cobble 128.1 - 256 mm			0				570				+	
Small Boulder 256.1 - 512 mm												
Medium Boulder 512.1 - 1024 mm												
Large Boulder 1024 mm & Larger												
TOTAL		0									The same	

2008 Burp

Page 4 of 8

80 00 00

SITE ID: 2 0

FORMATION CAPITAL CORPORATION, U.S.
IDAHO COBALT PROJECT
NPDES FISH TISSUE & MACROINVERTEBRATE SUMMARY REPORT

APPENDIX G

FISH TISSUE / METAL RESULTS

TABLE 4 Big Deer Creek -01
Metals Dry Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012

ounty, Idaho Lab ID	K1209055-031	K1209055-031D	K1209055-033	K1209055-035	K1209055-037	K1209055-039	K1209055-041	K1209055-043	K1209055-045	K1209055-047	K1209055-049	
Hildebrand Sample ID	BDC-01-01	BDC-01-01	BDC-01-02	BDC-01-03	BDC-01-04	BDC-01-05	BDC-01-06	BDC-01-07	BDC-01-08	BDC-01-09	BDC-01-10	Notes
Sample Date	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	
Sample Time	10:24	10:24	10:26	10:28	10:29	10:30	10:32	10:34	10:35	10:37	10:39	
RACE METALS - EPA METHOD 200.8												
Units 1	mg/Kg	тд/Кд	mg/Kg	mg/Kg 2.40	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Aluminum, Total	8.50	7.80	0.8 J		12.00	2.0 J	1.1 J	2.30	3.90	2.20	1.4 J	
Arsenic, Total	0.88	0.89	0.43 J	0.29 J	0.44 J	0.15 J	1.07	0.23 J	0.60	0.45 J	0.55	
Cadmium, Total	0.010 J	0.012 J	0.015 J	0.017 J	0.021	0.007 J	0.011 J	0.027	0.053	0.023	0.019 J	
Cobalt, Total	1.310	1.357	0.84	0.7360	0.414	0.645	0.454	1.040	0.557	0.901	1.070	
Lead. Total	0.0036 J	0.0043 J	0.0024 J	0.0046 J	0.0060 J	0.0040 J	0.0033 J	0.0045 J	0.0214	0.0166 J	0.0057 J	
Manganese, Total	1.04	1.05	0.96	1,6300	0.85	0.79	0.61	1.45	0.7600	0,7000	0.88	
Nickel, Total	0.08 J	0.10 J	0.02 J	0.02 U	0.08 J	0.02 J	0.02 U	0.26	0.02 U	0.02 J	0.02 U	
Thallium, Total	0.0167 J	0.0185 J	0.0160 J	0.0230	0.0140 J	0.0267	0.0088 J	0.0199	0.0153 J	0.0268	0.0044 J 32.0	
Zinc, Total	28.90	28.89	24.9	28.4	26.0	20.1	21.5	31.8	37.5	30.9	32.0	
EPA METHOD 1631E		A STATE OF THE STA		THE RESERVE OF THE PERSON NAMED IN								
Units ²	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	0.148	••	0.238	0.165	0.300	0.133	0.170	0.184	0.240	0.201	0.0774	ALCOHOL:
EPA METHOD 7742					38600 1 1 1 1 3	Marian Company				and the state of t		
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Selenium, Total	1.28 °	1.74 *	0.92 *	1.86 *	1.36 *	2.02 *	2.05 *	2.08 *	1.91 *	1.66 *	0.63 *	

- Metals results reported in milligram per kilogram
- Metals results reported in nanogram per gram
- Metals results reported in milligram per kilogram
- Total solids results reported in percent
- The result is an estimated value.

 The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL identification
- 1D

- Analyte present below calibration ratio
 The result is an outlier. See case narrative
 See mercury matrix spike/duplicate matrix spike summaries, pages 32-40

TABLE 5 Big Deer Creek - 01 Metals Wet Weight Basis
Formation Capitat Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhi, County, Idaho

Lab ID	K1209055-031	K1209055-031D	K1209055-033	K1209055-035	K1209055-037	K1209055-039	K1209055-041	K1209055-043	K1209055-045	K1209055-047	K1209055-049	
Hildebrand Sample ID	BDC-01-01	BDC-01-01	BDC-01-02	BDC-01-03	BDC-01-04	BDC-01-05	BDC-01-06	BDC-01-07	BDC-01-08	BDC-01-09	BDC-01-10	Notes
Sample Date	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Sample Time	10:24	10:24	10:26	10:28	10:29	10:30	10:32	10:34	10:35	10:37	10:39	
TRACE METALS - EPA METHOD 200.8	ENDERSON TRIPS									<u> </u>		
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Aluminum, Total	1.95	N/A	0.19 J	0.54	2.50	0.45 J	0.23 J	0.54	0.75	0.49	0.32 J	
Arsenic, Total	0.20	N/A	0.096 J	0.067 J	0.092 J	0.035 J	0.234	0.053 J	0.114	0.103 J	0.124	
Cadmlum, Total	0.0022 J	N/A	0.0034 J	0.0038 J	0.0044	0.0016 J	0.0023 J	0.0062	0.0101	0.0052	0.0043 J	
Cobalt, Total	0.302	N/A	0.19	0.168	0.0861	0.146	0.0995	0.24	0.106	0.205	0.240	
Lead, Total	0.0008 J	N/A	0.0005 J	0.0011 J	0.0013 J	0.0009 J	0.0007 J	0.0010 J	0.0041	0.0038 J	0.0013 J	
Manganese, Total	0.24	N/A	0.216	0.372	0.177	0.178	0.133	0.333	0.145	0.160	0.198	
Nickel, Total	0.019 J	N/A	0.005	0,004 U	0.016 J	0.005 J	0.004 U	0.060	0.004 U	0.005 J	0.004 U	
Thallium, Total	0,0038 J	N/A	0.0036	0.0048	0.0029 J	0.0061	0.0019 J	0.0046	0.0029	0.0061	0.0010J	
Zinc, Total	6.65	N/A	5.64	6.48	5.42	4.57	4.7	7.31	7.14	7.1	7.21	
EPA METHOD 1631E												
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	0.0340	**	0,0538	0.0376	0.0624	0.0302	0.0372	0.0423	0.0458	0.0458	0.0174	
EPA METHOD 7742												
Units ²	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Selenium, Total	0.29	N/A	0.21 *	0.42 °	0.28 *	0.46 *	0.45 *	0.48 *	0.36 *	0.38 *	0.14 *	
FREEZE DRY												
Units*	%	%	%	%	%	%	%	%	%	%	%	
Total Solids*	23	23.1	22.6	22.8	20.8	22.7	21.9	23	19.1	22.8	22.5	

- Metals results reported in milligram per kilogram
- Metals results reported in nanogram per gram
- Metals results reported in milligram per kilogram
- Total solids results reported in percent
- Total solids are only expressed on a wet weight basis
- The result is an estimated value.
- The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL identification
- Analyte present below calibration ratio N/A %
- Percent
- The result is an outlier. See case narrative
- See mercury matrix spike/duplicate matrix spike summaries, pages 32-40

TABLE 6
Big Deer Creek - 02
Metals Dry Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012

embi.	County.	Idaho	

County, Idaho Lab ID	K1209055-051	K1209055-053	K1209055-055	K1209055-057	K1209055-059	K1209055-061	K1209055-063	K1209055-065	K1209055-067	K1209055-069	
Hildebrand Sample ID	BDC-02-01	BDC-02-02	BDC-02-03	BDC-02-04	BDC-02-05	BDC-02-06	BDC-02-07	BDC-02-08	BDC-02-09	BDC-02-10	Notes
Sample Date	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	
Sample Time	12:39	12:41	12:43	12:45	12:46	12:48	12:50	12:51	12:52	12:54	
Outsiple 18800											
RACE METALS - EPA METHOD 200.8	BEST BANKS NAME										
Units'	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Aluminum, Total	4.20	1.0 J	1.0 J	1.1 J	1.5 J	141	1.7 J	1.2 J	1.3 J	1.0 J	
Arsenic, Total	0.40 J	1.05	0.15 J	0.49 J	0.63	0.62	1.24	0.34 J	0.85	0.76	
Cadmium, Total	0.058	0.026	0.027	0.030	0.031	0.051	0.068	0.062	0.026	0.061	
Cobalt, Total	0.116	0.035	0.061	0.080	0.156	0.539	0.081	0.068	0.037	0.086	
Lead, Total	0.0111 J	0.0015 J	0.0008 J	0.0161 J	0.0012 J	0.0475	0.0028 J	0,0015 J	0.0087 J	0.0104 J	
Manganese, Total	0.95	0.62	0.72	0.67	0.73	3.49	1.01	0.7100	0.6600	0.79	
Nickel, Total	0.02 U	0 03 J	0.02 U	0.1500	0.02 U	0.36	0.02 U	0.02 U	0.04 J	0.04 J	
Thailium, Total	0.0133 J	0.0150 J	0.0151 J	0.0207	0.0118 J	0.0183 J	0.0163 J	0.0144 J	0.0142 J	0.0175 J	
Zinc, Total	34.4	25.9	25.4	21.7	36.8	23.4	27.2	41.2	25.8	27.6	
EPA METHOD 1631E											
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	0.498	0.437	0.378	0.271	0.168	0.243	0.231	0.298	0.219	0.226	
EPA METHOD 7742						The state of the s					
Units'	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
	1,50 °	2.58 *	1.83°	1.51 °	1.99 °	1.73 *	0.29*	1.52 *	1.08 *,	2.66 *	
Selenium, Total	1,50 *	2.58 *	1.83*	1.51 *	1.99 °	1.73 *	0.29*	1.52	1.08	2.00	

S

- Metals results reported in milligram per kilogram Metals results reported in nanogram per gram
- Metals results reported in milligram per kilogram
- Total solids results reported in percent
 The result is an estimated value.
 The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL identification
- ID * The result is an outlier. See case narrative

TABLE 7
Big Deer Creek - 02
Metals Wet Weight Basis Idaho Cobalt Project & Lemhi, County, Idaho

rmation Capital Corporation		
sho Cobalt Project Monitoring Summary, Sentember, 2012		

Lab ID	K1209055-051	K1209055-053	K1209055-055	K1209055-057	K1209055-059	K1209055-061	K1209055-063	K1209055-065	K1209055-067	K1209055-069	
Hildebrand Sample ID	BDC-02-01	BDC-02-02	BDC-02-03	BDC-02-04	BDC-02-05	BDC-02-06	BDC-02-07	BDC-02-08	BDC-02-09	BDC-02-10	Notes
Sample Date	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	9/4/2012	Hotes
Sample Time	12:39	12:41	12:43	12:45	12:46	12:48	12:50	12:51	12:52	12:54	
RACE METALS - EPA METHOD 200.8			Chicagons - Harasin				nja sistanjanjanjan ija				
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Aluminum, Total	0,87	0.21 J	0.23 J	0.24 J	. 0.31 J	28.4	0.38 J	0.24 J	0.27 J	0.22 J	
Arsenic, Total	0.82 J	0.223	0.035 J	0.108 J	0.132	0.124	0.272	0.071 J	0.186	0.163	
Cadmium, Total	0.0120	0.0056	0.0063	0.0067	0.0066	0.0102	0.0149	0.0128	0.0057	0.0131	
Cobalt, Total	0.0239	0.0074	0.0140	0.0177	0.0325	0,108	0.0179	0.0141	0.0080	0.0186	
Lead, Total	0.0023 J	0.0003 J	0.0002 J	0.0036 J	0.0003 J	0.0095	0.0006 J	0.0003 J	0.0019 J	0.0022 J	
Manganese, Total	0.195	0.131	0.166	0.149	0.153	0.702	0.222	0.147	0.144	0.171	
Nickel, Total	0.004 U	0.006 J	0.004 U	0.032 J	0.004 U	0.073	0.004 U	0.004 U	L 600'0	0.008 J	
Thallium, Total	0.0027 J	0.0032 J	0.0035 J	0.0046	0.0025 J	0.0037	0.0036 J	0.0030 J	0.0031 J	0.0038 J	
Zinc, Total	7.08	5.52	5,85	4.64	7.70	4.71	5,98	8.54	5.63	5.96	
EPA METHOD 1631E			A STATE OF STATE								
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	0.103	0.0931	0.0869	0.0604	0.0347	0.0491	0.0508	0.0617	0.0477	0.0488	
EPA METHOD 7742											Service Contract
Units'	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Selenium, Total	0.31 *	0.55 *	0.42*	0.34 *	0.42 *	0.35 *	0.72 *	0.32 *	0.24 *	0.57 *	
FREEZE DRY	11 10 10 10										
Units*	%	%	%	%	%	%	%	%	%	%	
Total Solids	20.6	21.3	23	22.3	20.9	20.2	22	20.7	21.8	21.6	

- Metals results reported in milligram per kilogram
- Metals results reported in minigram per Miogram

 Metals results reported in milligram per kilogram

 Total solids results reported in percent
- Total solids are only expressed on a wet weight basis
- J The result is an estimated value.
 U The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL ID Identification
- U 1D %
- Percent
 The result is an outlier. See case parrative

TABLE 8
Upper Blackbird Creek - 01
Metals Dry Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhi, County, Idaho

, County, IdahoLab ID	K1209055-071	K1209055-071D	K1209055-073	K1209055-075	K1209055-077	K1209055-079	K1209055-081	K1209055-081D	K1209055-083	K1209055-085	K1209055-087	K1209055-089	
Hildebrand Samole ID	UBC-01	UBC-01	UBC-02	UBC-03	UBC-04	UBC-05	UBC-06	UBC-06	UBC-07	UBC-08	UBC-09	UBC-10	Notes
Sample Date	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	
Sample Time	10:24	10:24	10:26	10:28	10:29	10:30	10:32	10:32	10:34	10:35	10:37	10:39	
RACE METALS - EPA METHOD 200.8	9-10-11												
Units'	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	រាជ្ជ/Kg	
Aluminum, Total	5.00	6.60	41.90	2.90	1.0 J	1.1 J	2.1	N/A	0.5 J	2.00	0.6 J	2.10	Chain of custody has sample I
Arsenic, Total	0.85	0.86	0.97	0.32 J	1.26	0.63	0.14 J	N/A	0.68	0.63	0.54	0.26 J	for Upper Black Bird Creek as
Cadmium, Total	0.015 J	0.016 J	0.011 J	0.019 J	0.018 J	0.011 J	0.015 J	N/A	0.010 J	0.037	0.011 J	0.029	UBC, however, laboratory reporte
Cobalt, Total	1,430	1,455	0.908	0.565	1.92	1.11	0.202	N/A	0.955	0.951	0.635	0.491	as MBC
Lead, Total	N/A	N/A	0.0057 J	0.0036 J	0.0014 J	0.0017 J	0.0025 J	0.0641	0.0041 J	0.0016 J	0.0011 J	0.0016 J	
Manganese, Total	4.51	4.59	13.90	2.47	18.10	5.81	0.64	N/A	5.98	6.25	3.09	1,08	
Nickel, Total	0.13 J	0.13 J	0.11 J	0.03 J	0.08 J	0.04 J	0.03 J	N/A	0.03 J	0.03 J	0.02 J	0.03 J	
Thallium, Total	0.0097 J	0.01183	0.0019 J	0.0025 J	0.0077 J	0.0064 J	0.0040 J	N/A	0.0088	0.0060 J	0.0027 J	0.0064 J	
Zinc, Total	58.42°	77,10 *	24.8 °	35.5 °	34.5 *	30.3 °	34.1 *	N/A	42.9 *	43.1 *	44.1 °	40.0 *	
EPA METHOD 1631E						A STATE OF THE STA							
Units ²	mo/Ka	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	. 0.618	**	0.211	0.242	0.181	0.161	0.180	**	0.154	0.326	0.220	0.294	
EPA METHOD 7742		Environment of the second				THE RESERVE TO SERVE							
Units*	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	т9/Кр	
Selenium, Total	1.40	0.98	0.51	1.04	1.32	1.65	1.20	N/A	1.93	1.40	1 85	1.86	
	ELCS COLUMN TO BE SELECTED.	CHALL CONTRACTOR											

- fotes:

 Metals results reported in milligram per kilogram

 Metals results reported in nanogram per gram

 Metals results reported in milligram per kilogram

 Total solids results reported in percent

 The result is an estimated value.

 ID Identification

 Analyte present below calibration ratio

 N/A Not available

 See mercury matrix spike/duplicate matrix spike summaries, pages 32-40

 The result is an outlier. See case narrative

TABLE 9

TABLE 9
Upper Blackbird Creek - 01
Metals Wet Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lembi, County, Idaho

County, idaho Lab ID	K1209055-071	K1209055-071D	K1209055-073	K1209055-075	K1209055-077	K1209055-079	K1209055-081	K1209055-081D	K1209055-083	K1209055-085	K1209055-087	K1209055-089	
Hildebrand Sample ID	UBC-01	UBC-01	UBC-02	UBC-03	UBC-04	UBC-05	UBC-06	UBC-06	UBC-07	UBC-08	UBC-09	UBC-10	Notes
Sample Date	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	9/5/2012	110103
Sample Time	10:24	10:24	10:25	10:28	10:29	10:30	10:32	10:32	10:34	10:35	10:37	10:39	<u> </u>
FRACE METALS - EPA METHOD 200.8					Mary and							60.	
Units'	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	1
Aluminum, Total	0.94	N/A	10.2	0.59	0.21 J	0.23 J	0.45	N/A	0.11 J	0.42	0.12 J	0.43	Chain of custody has sample
Arsenic, Total	0.160	N/A	0.236	0.065 J	0.267	0.130	0.030 J	N/A	0.140	0.130	0.109	0.054 J	for Upper Black Bird Creek a
Cadmium, Total	0.0028 J	N/A	0.0028 J	0,0039 J	0.0039 J	0.0024 J	0.0033 J	N/A	0.0021 J	0.0076	0.0023 J	0.0060	UBC, however, laboratory report
Cobalt, Total	0.271	N/A	0.221	0.177	0,406	0.229	0.0430	N/A	0.197	0.197	0.129	0.100	as MBC
Lead. Total	0.0006 J	N/A	0.0014 J	0.0007 J	0.0003 J	0.0004 J	0.0005 J	N/A	0.0008 J	0.0003 J	0.0002 J	0.0003 J	
Manganese, Total	0.853	N/A	3.380	0.511	3.830	1.200	0.137	N/A	1.230	1.290	0.628	0.221	
Nickel, Total	0.025 J	N/A	0.027 J	0.007 J	0.017 J	0.009 J	0.006 J	N/A	0.005 J	0.006 J	0.004 J	0.005 J	
Thallium, Total	0.0018 J	N/A	0.0005 J	0.0005 J	0.0016 J	0.0013 J	0.0008 J	N/A	0.0018 J	0.0012 J	0.0005 J	0.0013 J	
Zinc, Total	11.0 *	N/A	6.02 °	7,35 *	7.30 *	6.28 *	7.26 °	N/A	8.83 *	8.92 *	8.96 °	8.16 *	
EPA METHOD 1631E	Texas ()						District of	The Color of the Color					THE STATE OF THE S
Units*	mg/Kg	mg/Kg	mg/Kg	mq/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	0.117	**	0.0513	0.0501	0.0384	0.0333	0.0383	69	0.0317	0.0675	Ü.0447	0.0600	
EPA METHOD 7742	and the positions		THE PARTY NAMED IN										
Units	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	mg/Kg	
Selenium, Total	0.26	N/A	0.12	0.22	0 28	0.34	0.25	N/A	0.40	0.29	0.38	0.38	
FREEZE DRY				4					y Project of				
Units	%	%	%	1 %	%	1 %	%	%	%	%	1 %	%	
Total Solids*	19	19.3	24.3	20.7	21,2	20.7	21.3	N/A	20.6	20.7	20.3	20.4	

- Metals results reported in milligram per tidlogræm Metals results reported in nanogram per क्रका
- Metals results reported in milligram per kilogram
- Total solids results reported in mingratin be actional.

 Total solids are only expressed on a wet weight basis

 The result is an estimated value.

 Identification

- J Identification
 J Analyte present below calibration ratio
 Percent
 N/A Not available
 See mercury matrix spike/duplicate matrix spike summaries, pages 32-40
 The result is an outlier. See case narrative

TABLE 10
Method Blanks Dry Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhi, County, Idaho

Lab ID	K1209055-MB1	K1209055-MB2	
Hildebrand Sample ID	N/A	N/A	Notes
Sample Date	10/8/2012	10/8/2012	
TRACE METALS - EPA METHOD 200.8		0.50	
Units'	mg/Kg	mg/Kg	E4
Aluminum, Total	0.2 U	0.4 J	
Arsenic, Total	0.02 U	0.02 U	
Cadmium, Total	0.002 U	0.002 U	- 1
Cobalt, Total	0.003 U	0.003 U	41
Lead, Total	0.0009 J	0.0015 J	Analyzed on 10/15/12
Manganese, Total	0.02 U	0.02 U	P
Nickel, Total	0.02 U	0.02 U	
Thallium, Total	0.0027 J	0.0009 U	
Zinc, Total	0.08 J	0.38 J *	
EPA METHOD 7742			
Units	mg/Kg	mg/Kg	
Selenium, Total	0.05 * U	0.05 U	Analyzed on 10/10/12

- Metals results reported in milligram per kilogram
- J The result is an estimated value.
- ID Identification

 * The result is a
- * The result is an outlier. See case narrative.
- U The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL.

TABLE 11
Method Blanks Wet Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhi, County, Idaho

Lab ID	K1209055-MB1	K1209055-MB2	
Hildebrand Sample ID	N/A	N/A	Notes
Sample Date	10/8/2012	10/8/2012	
TRACE METALS - EPA METHOD 200.8			
Units'	mg/Kg	mg/Kg	
Aluminum, Total	0.04 U	0.08 J	
Arsenic, Total	0.004 U	0.004 U	
Cadmium, Total	0.0004 U	0.0004 U	
Cobalt, Total	0.0006 U	0.0006 U	
Lead, Total	0.0002 J	0.0003 J	Analyzed on 10/15/12
Manganese, Total	0.004 U	0.004 U	
Nickel, Total	0.004 U	0.004 U	
Thallium, Total	0.0005 J	0.0002 U	
Zinc, Total	0.02 J	0.38 J *	
EPA METHOD 7742			
Units'	mg/Kg	mg/Kg	
Selenium, Total	0.01 * U	0.01 U	Analyzed on 10/10/12

- Metals results reported in milligram per kilogram
- J The result is an estimated value.
- ID Identification
- * The result is an outlier. See case narrative.
- U The analyte was analyzed for, but was not detected ("Non-detect") at or above the MRL/MDL.

TABLE 12
Mercury Method Blanks Dry Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhi, County, Idaho

Lab ID Hildebrand Sample ID	K1209055-MB1 N/A	K1209055-MB2 N/A	K1209055-MB3 N/A	Notes
Sample Date	10/19/2012	10/19/2012	10/19/2012	
EPA METHOD 1631E				
Units	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	ND	ND.	ND ND	

- Metals results reported in nanogram per gram
- J The result is an estimated value.
- ID Identification
- * The result is an outlier. See case narrative.
- ND Not Detected

TABLE 13
Mercury Method Blanks Wet Weight Basis
Formation Capital Corporation
Idaho Cobalt Project Monitoring Summary, September, 2012
Lemhl, County, Idaho

Lab ID	K1209055-MB1	K1209055-MB2	K1209055-MB3	
Hildebrand Sample ID	N/A	N/A	N/A	Notes
Sample Date	10/19/2012	10/19/2012	10/19/2012	
EPA METHOD 1631E				
	Sandal Control of			
Units'	mg/Kg	mg/Kg	mg/Kg	
Mercury, Total	ND ND	ND	ND	

Metals results reported in nanogram per gram
J The result is an estimated value.
ID identification
The result is an outlier. See case narrative.
ND Not Detected